

## Chapter Four: Real functions with real variable

### 4.1 Generalities

#### Definition 4.1

We call a real function of a real variable every application  $f$  of a subset  $D$  of  $\mathbb{R}$  on set  $\mathbb{R}$ .

$D$  is called the domain of definition for  $f$ .

We call the graph of the function  $f$  the subset of  $\mathbb{R}^2$  which we denote by  $\Gamma_f$ , and defined as follows:  $\Gamma_f = \{(x; y) \in \mathbb{R}^2; x \in D \wedge y = f(x)\}$  or  $\Gamma_f = \{(x; f(x)); x \in D\}$ .

The image of the domain  $D$  by  $f$  is denoted by  $f(D)$  where:  $f(D) = \{y \in \mathbb{R}; \exists x \in D: y = f(x)\}$ .

Definition 4.2 Let  $f: D \rightarrow \mathbb{R}$  be a function.

We say that the function  $f$  is bounded from above (bounded from below, respectively) if, and only if, the set  $f(D)$  is bounded from above (bounded from below, respectively)

So, ( $f$  is bounded from above)  $\Leftrightarrow (\exists M \in \mathbb{R}; \forall x \in D: f(x) \leq M)$ .

, ( $f$  is bounded from below)  $\Leftrightarrow (\exists m \in \mathbb{R}; \forall x \in D: f(x) \geq m)$ .

We say that the function  $f$  is bounded if, and only if, it is bounded from above and from below.

So, ( $f$  is bounded)  $\Leftrightarrow (\exists M \in \mathbb{R}_+^*; \forall x \in D: |f(x)| \leq M)$ .

#### Remark 4.1

If the function  $f$  is bounded on  $D$ , then the part  $f(D)$  is bounded on  $\mathbb{R}$ . It accepts an upper bound and a lower bound, which we denote by  $Sup_D f$  and  $Inf_D f$  respectively.

Definition 4.3 Let  $f: D \rightarrow \mathbb{R}$  be a function.

We say that  $f$  is increasing over  $D$  (strictly increasing, respectively) if and only if

$\forall x; y \in D: x < y \Rightarrow f(x) \leq f(y)$  ( $\forall x; y \in D: x < y \Rightarrow f(x) < f(y)$ , respectively).

We say that  $f$  is decreasing over  $D$  (strictly decreasing, respectively) if and only if

$\forall x; y \in D: x < y \Rightarrow f(x) \geq f(y)$  ( $\forall x; y \in D: x < y \Rightarrow f(x) > f(y)$ , respectively).

We say that  $f$  is constant over  $D$  if and only if  $\forall x; y \in D: x \neq y \Rightarrow f(x) = f(y)$ .

**Definition 4.4** Let  $f: D \rightarrow \mathbb{R}$  be a function.

We say that  $f$  have a local maximum (local minimum, respectively) at point  $x_0$  of  $D$  if:

$\exists \alpha \in \mathbb{R}_+^*; \forall x \in D: |x - x_0| < \alpha \Rightarrow f(x) \leq f(x_0)$  ( $f(x) \geq f(x_0)$ ), respectively).

And if  $\forall x \in D: f(x) \leq f(x_0)$  ( $f(x) \geq f(x_0)$ ), respectively) we say that  $f$  have an absolute maximum (absolute minimum, respectively) at  $x_0$ .

## **4.2 limit of a function**

### **4.2.1 Finite limit**

#### **Definition 4.5**

We say a subset of  $\mathbb{R}$  is a <sup>جوار</sup>neighborhood for a point  $x_0$  of  $\mathbb{R}$  if it contains an open interval that includes  $x_0$ . And we symbolize it with  $V_{x_0}$ .

Let  $f$  be a function, defined on a neighborhood  $V_{x_0}$  of point  $x_0$ .

We say that the function  $f$  has a limit  $\ell$  ( $\ell \in \mathbb{R}$ ) at point  $x_0$  if, and only if,

$\forall \varepsilon > 0; \exists \delta > 0; \forall x \in V_{x_0}: 0 < |x - x_0| < \delta \Rightarrow |f(x) - \ell| < \varepsilon$ , we write  $\lim_{x \rightarrow x_0} f(x) = \ell$ .

#### **Remark**

We say that  $f$  does not accept the number  $\ell$  as a limit at  $x_0$  if and only if

$$\exists \varepsilon > 0; \forall \delta > 0; \exists x \in V_{x_0}: 0 < |x - x_0| < \delta \text{ و } |f(x) - \ell| \geq \varepsilon$$

#### **proposition 4.1**

If  $\lim_{x \rightarrow x_0} f(x) = \ell \neq 0$ , then there exists a domain of the form  $]x_0 - \alpha, x_0[ \cup ]x_0, x_0 + \alpha[$ , with  $\alpha > 0$ ,

such that  $f(x)$  has the same sign as  $\ell$ .

#### **Proof**

For  $\varepsilon = |\ell|$ , then  $\exists \alpha > 0; \forall x \in V_{x_0}: 0 < |x - x_0| < \alpha \Rightarrow |f(x) - \ell| < |\ell|$  from him

$$x \in ]x_0 - \alpha, x_0[ \cup ]x_0, x_0 + \alpha[ \Rightarrow \begin{cases} 2\ell < f(x) < 0; & \ell < 0 \\ 0 < f(x) < 2\ell; & \ell > 0 \end{cases}$$

$\Rightarrow f(x)$  has the same sign as  $\ell$ .

### Examples

1) Let  $f: x \rightarrow 5x - 7$  Be a function , using the definition prove that:  $\lim_{x \rightarrow 2} f(x) = 3$ .

Since  $f$  is defined on  $\mathbb{R}$ , we can take  $V_2 = \mathbb{R}$ . ( $V_2$  is a neighborhood of point 2 )

Let  $\varepsilon \in \mathbb{R}_+^*$ , we have  $\forall x \in \mathbb{R}$ :

$$\begin{aligned} |f(x) - 3| < \varepsilon &\Leftrightarrow |5x - 7 - 3| < \varepsilon \\ &\Leftrightarrow |x - 2| < \frac{\varepsilon}{5} \end{aligned}$$

So it is enough to take  $\delta = \frac{\varepsilon}{5}$  to achieve the following:

$$\forall \varepsilon > 0; \exists \delta > 0; \forall x \in \mathbb{R} : 0 < |x - 2| < \delta \Rightarrow |f(x) - 3| < \varepsilon.$$

2) Let  $f: x \rightarrow \frac{1}{x+1}$  Be a function , using the definition prove that:  $\lim_{x \rightarrow 1} f(x) = \frac{1}{2}$ .

Since  $f$  is defined on  $\mathbb{R} - \{-1\}$ , we can take  $V_1 = [0; +\infty[$ . ( $V_1$  is a neighborhood of point 2 )

Let  $\varepsilon \in \mathbb{R}_+^*$ , we have

$$\forall x \in V_1: \left| f(x) - \frac{1}{2} \right| = \left| \frac{1}{x+1} - \frac{1}{2} \right| = \frac{|x-1|}{2|x+1|} < \frac{|x-1|}{2}.$$

Therefore, it suffices to take  $\frac{|x-1|}{2} < \varepsilon$  to be  $\left| f(x) - \frac{1}{2} \right| < \varepsilon$ , from which

$$\left| \frac{x-1}{2} \right| < \varepsilon \Leftrightarrow |x - 1| < 2\varepsilon. \text{ So it is enough to take } \delta = 2\varepsilon \text{ to achieve the following:}$$

$$\forall \varepsilon > 0; \exists \delta > 0; \forall x \in V_1: 0 < |x - 1| < \delta \Rightarrow \left| f(x) - \frac{1}{2} \right| < \varepsilon.$$

### Definition 4 6

Let  $f$  be a function defined in the interval  $V_{x_0} = ]x_0, b[$ , we say that  $f$  have the limit  $\ell$  from the right at  $x_0$  if and only if

$$\forall \varepsilon > 0; \exists \delta > 0; \forall x \in V_{x_0}: 0 < x - x_0 < \delta \Rightarrow |f(x) - \ell| < \varepsilon.$$

we write  $\lim_{x \rightarrow x_0^+} f(x) = \ell$  or  $\lim_{x \rightarrow x_0^+} f(x) = \ell$ .

Let  $f$  be a function defined in the interval  $V_{x_0} = ]a, x_0[$ , we say that  $f$  have the limit  $\ell$  from the left at  $x_0$  if and only if

$$\forall \varepsilon > 0; \exists \delta > 0; \forall x \in V_{x_0}: -\delta < x - x_0 < 0 \Rightarrow |f(x) - \ell| < \varepsilon.$$

we write  $\lim_{x \rightarrow x_0^-} f(x) = \ell$  or  $\lim_{x \rightarrow x_0^-} f(x) = \ell$ .

### Proposition 4.2

A function  $f$  has a limit at  $x_0$  if and only if it accepts right and left limits at  $x_0$  and these limits are equal.

### Example

Let the function  $f$  defined on  $\mathbb{R}$  by  $f(x) = \begin{cases} 3x - 1 & \text{if } x \leq 1 \\ \frac{6}{x+2} & \text{if } x > 1 \end{cases}$ .

Prove that:  $\lim_{x \rightarrow 1^+} f(x) = 2$  and  $\lim_{x \rightarrow 1^-} f(x) = 2$  what do you conclude.

1) Let  $V_1 = ]-\infty; 1]$  and  $\varepsilon \in \mathbb{R}_+^*$ , we have

$$\begin{aligned} \forall x \in V_1: |f(x) - 2| < \varepsilon &\Leftrightarrow |3x - 3| < \varepsilon \\ |3x - 3| < \varepsilon &\Leftrightarrow 0 < |x - 1| < \frac{\varepsilon}{3} \\ &\Leftrightarrow 0 < -x + 1 < \frac{\varepsilon}{3} \\ &\Leftrightarrow -\frac{\varepsilon}{3} < x - 1 < 0 \end{aligned}$$

It is enough to take  $\delta = \frac{\varepsilon}{3}$  to achieve the following:

$$\forall \varepsilon > 0; \exists \delta > 0; \forall x \in V_1: 0 < 1 - x < \delta \Rightarrow |f(x) - 2| < \varepsilon$$

Let  $V_1 = [1; +\infty[$  and  $\varepsilon \in \mathbb{R}_+^*$ , we have

$$\forall x \in V_1: |f(x) - 2| = \frac{2|x - 1|}{x + 2} < \frac{2}{3}|x - 1|$$

So

$$\frac{2}{3}|x - 1| < \varepsilon \Leftrightarrow |x - 1| < \frac{3}{2}\varepsilon \Leftrightarrow 0 < x - 1 < \frac{3}{2}\varepsilon$$

It is enough to take  $\delta = \frac{3\varepsilon}{2}$  to achieve the following:

$$\forall \varepsilon > 0; \exists \delta > 0; \forall x \in V_1: 0 < x - 1 < \delta \Rightarrow |f(x) - 2| < \varepsilon$$

**Conclusion:** Since  $\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = 2$   $f$  accepts a limit at 1, which is 2.

### Theorem 4.1

If a function  $f$  accepts a limit at  $x_0$ , then this limit is unique.

### Proof

Let  $f$  accept two different limits  $\ell$  and  $\ell'$  where  $\ell > \ell'$ .

for  $\varepsilon = \frac{\ell - \ell'}{2}$ ;  $\exists \delta_1, \delta_2 > 0$ ;  $\forall x \in V_{x_0}$ :

$$0 < |x - x_0| < \delta_1 \Rightarrow |f(x) - \ell| < \varepsilon = \frac{\ell - \ell'}{2}$$

and

$$0 < |x - x_0| < \delta_2 \Rightarrow |f(x) - \ell'| < \varepsilon = \frac{\ell - \ell'}{2}$$

For  $\delta = \min\{\delta_1, \delta_2\}$  Then  $\forall x \in V_{x_0}$ :

$$\begin{aligned} 0 < |x - x_0| < \delta &\Rightarrow |\ell - \ell'| = |f(x) - \ell - (f(x) - \ell')| \\ &\Rightarrow |\ell - \ell'| < \varepsilon + \varepsilon = 2\varepsilon \\ &\Rightarrow |\ell - \ell'| < |\ell - \ell'| \end{aligned}$$

This is a contradiction. So  $\ell = \ell'$

#### **4.2.2 Limit of a function using sequences**

##### **Theorem 4.2**

Let  $f: D \rightarrow \mathbb{R}$  be a function and  $x_0 \in D$ . The following two conditions are equivalent.

1)  $\lim_{x \rightarrow x_0} f(x) = \ell$ .

2) For all sequence  $(x_n)$  where  $\forall n \in \mathbb{N}: x_n \in D \wedge x_n \neq x_0$  then:

$$\left(\lim_{n \rightarrow +\infty} x_n = x_0\right) \Rightarrow \left(\lim_{n \rightarrow +\infty} f(x_n) = \ell\right)$$

##### **Proof**

##### **Necessary condition:**

We impose  $\lim_{x \rightarrow x_0} f(x) = \ell$  and let  $(x_n)$  sequence where  $\forall n \in \mathbb{N}: x_n \in D \wedge x_n \neq x_0$  and  $\lim_{n \rightarrow \infty} x_n =$

$x_0$ . Let us prove that:  $\lim_{n \rightarrow +\infty} f(x_n) = \ell$ .

For  $\varepsilon > 0$  then  $\exists \delta > 0$ ;  $\forall x \in V_{x_0}: 0 < |x - x_0| < \delta \Rightarrow |f(x) - \ell| < \varepsilon$ . So

$\exists N \in \mathbb{N}; \forall n \in \mathbb{N}: n > N \Rightarrow |x_n - x_0| < \delta \Rightarrow |f(x_n) - \ell| < \varepsilon$ .

So  $\forall \varepsilon > 0; \exists N \in \mathbb{N}; \forall n \in \mathbb{N}: n > N \Rightarrow |f(x_n) - \ell| < \varepsilon$ . So  $\lim_{n \rightarrow +\infty} f(x_n) = \ell$ .

**Sufficient condition:** We now assume that for every sequence  $(x_n)$  where

$$\forall n \in \mathbb{N}: x_n \in D \wedge x_n \neq x_0 \text{ then } (\lim_{n \rightarrow +\infty} x_n = x_0) \Rightarrow (\lim_{n \rightarrow +\infty} f(x_n) = \ell).$$

Let us prove that  $\lim_{x \rightarrow x_0} f(x) = \ell$ . Assume that  $\lim_{x \rightarrow x_0} f(x) \neq \ell$  that is

$$\exists \varepsilon > 0; \forall \delta > 0; \exists x \in V_{x_0}: 0 < |x - x_0| < \delta \text{ and } |f(x) - \ell| \geq \varepsilon.$$

and for  $\delta = \frac{1}{n}$  then  $\forall n \in \mathbb{N}^*; \exists x_n \neq x_0$  and  $x_n \in V_{x_0}: |x_n - x_0| < \frac{1}{n}$  and  $|f(x_n) - \ell| \geq \varepsilon$ .

So  $\lim_{n \rightarrow +\infty} x_n = x_0$  and  $\lim_{n \rightarrow +\infty} f(x_n) \neq \ell$  ( this is a contradiction ).

### **Remark**

To prove that a function  $f$  has no limit at  $x_0$ , it is enough to find two sequences  $(x_n)$  and  $(x'_n)$  that converge towards  $x_0$  but  $\lim_{n \rightarrow \infty} f(x'_n) \neq \lim_{n \rightarrow \infty} f(x_n)$  Or we are looking for a sequence  $(x_n)$  that converges toward  $x_0$  but the sequence  $(f(x_n))_{n \in \mathbb{N}}$  diverges.

### **Example**

Prove that the function  $f: x \rightarrow \cos \frac{1}{x}$  does not accept a limit at 0.

Let the sequences  $(x_n)$  and  $(x'_n)$  where  $\forall n \in \mathbb{N}^*: x_n = \frac{1}{2\pi n + \frac{\pi}{2}}, x'_n = \frac{1}{2\pi n + \pi}$ .

We have  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} x'_n = 0$  On the other hand:  $\forall n \in \mathbb{N}^*: f(x'_n) = -1; f(x_n) = 0$ .

So

$\lim f(x'_n) = -1; \lim f(x_n) = 0$  So:  $\lim f(x'_n) \neq \lim f(x_n)$  i.e.  $f$  does not accept a limit at 0.

### **4.2.3 Infinite limits**

We say a subset of  $\mathbb{R}$  is a <sup>جوار</sup>neighborhood of  $+\infty$  ( $-\infty$ , respectively) if it contains an open interval of the form  $]a, +\infty[$  ( $]-\infty, b[$ , respectively) And we symbolize it with  $V_{+\infty}$  ( $V_{-\infty}$ , respectively).

### **Definitions**

$$(\forall \varepsilon > 0; \exists A > 0; \forall x \in V_{+\infty}: x > A \Rightarrow |f(x) - \ell| < \varepsilon) \Leftrightarrow (\lim_{x \rightarrow +\infty} f(x) = \ell)$$

$$(\forall \varepsilon > 0; \exists A > 0; \forall x \in V_{-\infty}: x < -A \Rightarrow |f(x) - \ell| < \varepsilon) \Leftrightarrow (\lim_{x \rightarrow -\infty} f(x) = \ell)$$

$$(\forall A > 0; \exists \delta > 0; \forall x \in V_{x_0}: |x - x_0| < \delta \Rightarrow f(x) > A) \Leftrightarrow (\lim_{x \rightarrow x_0} f(x) = +\infty)$$

$$(\forall A > 0; \exists \delta > 0; \forall x \in V_{x_0}: |x - x_0| < \delta \Rightarrow f(x) < -A) \Leftrightarrow (\lim_{x \rightarrow x_0} f(x) = -\infty)$$

$$(\forall A > 0; \exists B > 0; \forall x \in V_{+\infty}: x > B \Rightarrow f(x) > A) \Leftrightarrow (\lim_{x \rightarrow +\infty} f(x) = +\infty)$$

$$(\forall A > 0; \exists B > 0; \forall x \in V_{+\infty}: x > B \Rightarrow f(x) < -A) \Leftrightarrow (\lim_{x \rightarrow +\infty} f(x) = -\infty)$$

$$(\forall A > 0; \exists B > 0; \forall x \in V_{-\infty}: x < -B \Rightarrow f(x) > A) \Leftrightarrow (\lim_{x \rightarrow -\infty} f(x) = +\infty)$$

$$(\forall A > 0; \exists B > 0; \forall x \in V_{-\infty}: x < -B \Rightarrow f(x) < -A) \Leftrightarrow (\lim_{x \rightarrow -\infty} f(x) = -\infty)$$

### Examples

1) Prove that  $\lim_{x \rightarrow \infty} \frac{2x}{x-1} = 2$ .

The function  $x \rightarrow \frac{2x}{x-1}$  is defined on  $V_{+\infty} = ]1; +\infty[$ , for  $\varepsilon \in \mathbb{R}_+$  we have

$$\forall x \in V_{+\infty}: |f(x) - 2| < \varepsilon \Leftrightarrow \frac{2}{|x-1|} < \varepsilon \Leftrightarrow \frac{2}{x-1} < \varepsilon \Leftrightarrow x > \frac{2}{\varepsilon} + 1$$

Therefore, it is sufficient to choose  $B = \frac{2}{\varepsilon} + 1$  to obtain:

$$\forall \varepsilon > 0; \exists B \in \mathbb{R}_+^*; \forall x \in V_{+\infty}: x > B \Rightarrow |f(x) - 2| < \varepsilon$$

2) Prove that  $\lim_{\substack{x \rightarrow 1 \\ x < 1}} \frac{2x}{x-1} = -\infty$ .

Let  $V_1 = ]0; 1[$ , for  $A \in \mathbb{R}_+^*$  we have

$$\begin{aligned} \forall x \in V_1: f(x) < -A &\Leftrightarrow \frac{2x}{x-1} < -A \Leftrightarrow 2 + \frac{2}{x-1} < -A \\ &\Leftrightarrow 0 > x-1 > \frac{2}{-A-2} \\ &\Leftrightarrow -\frac{2}{A+2} < x-1 < 0 \end{aligned}$$

Therefore, it is sufficient to choose  $\delta = \frac{2}{A+2}$  to obtain:

$$\forall A > 0; \exists \delta \in \mathbb{R}_+^*; \forall x \in V_1: 0 < 1-x < \delta \Rightarrow f(x) < -A.$$

### 4.2.4 Operation on limits

#### Theorem 4.3

Let  $f$  and  $g$  be functions defined on the neighborhood  $V_{x_0}$ , with the possible exception of  $x_0$ , where

$$\forall x \in V_{x_0}: f(x) < g(x)$$

1) If  $\lim_{x \rightarrow x_0} f(x) = \ell$  and  $\lim_{x \rightarrow x_0} g(x) = \ell'$  then  $\ell \leq \ell'$ .

2) If  $\lim_{x \rightarrow x_0} f(x) = +\infty$  then  $\lim_{x \rightarrow x_0} g(x) = +\infty$ .

3)  $\lim_{x \rightarrow x_0} g(x) = -\infty$  then  $\lim_{x \rightarrow x_0} f(x) = -\infty$ .

Let  $f, g$  and  $h$  be functions defined on the neighborhood  $V_{x_0}$ , with the possible exception of  $x_0$ , where  $\forall x \in V_{x_0}: h(x) < f(x) < g(x)$  and  $\lim_{x \rightarrow x_0} g(x) = \lim_{x \rightarrow x_0} h(x) = \ell$ , then

$$\lim_{x \rightarrow x_0} f(x) = \ell.$$

**Proof**

Assume that  $\forall x \in V_{x_0}: f(x) < g(x)$  and  $\lim_{x \rightarrow x_0} f(x) = \ell$ ,  $\lim_{x \rightarrow x_0} g(x) = \ell'$  and suppose that

$\ell > \ell'$ . For  $\varepsilon = \frac{\ell - \ell'}{2}$  then

$$\exists \delta_1 > 0: 0 < |x - x_0| < \delta_1 \Rightarrow |f(x) - \ell| < \varepsilon \Rightarrow \frac{\ell + \ell'}{2} < f(x) < \frac{3\ell - \ell'}{2}$$

$$\exists \delta_2 > 0: 0 < |x - x_0| < \delta_2 \Rightarrow |g(x) - \ell'| < \varepsilon \Rightarrow \frac{3\ell' - \ell}{2} < g(x) < \frac{\ell + \ell'}{2}$$

But taking  $\delta = \min\{\delta_1, \delta_2\}$  then  $0 < |x - x_0| < \delta \Rightarrow g(x) < \frac{\ell + \ell'}{2} < f(x)$  this is contradiction the hypothesis.  $\forall x \in V_{x_0}: f(x) < g(x)$ .

**Theorem 4.4**

If  $f$  and  $g$  are functions defined in the neighborhood  $V_{x_0}$ , with the possible exception of  $x_0$ , and have the limits  $\ell, \ell'$ , at  $x_0$  respectively, then the functions  $f + g, f - g, \lambda f, |f|$  it has the limits  $\ell + \ell', \ell - \ell', \lambda\ell, \ell\ell', |\ell|$ , at  $x_0$  respectively. And if  $\ell' \neq 0$ , then the function  $\frac{1}{g}$  it has the limit  $\frac{1}{\ell'}$  at  $x_0$ .

**Proof** (Let us prove the last case )

Assume that  $\lim_{x \rightarrow x_0} g(x) = \ell' \neq 0$  for  $\varepsilon = \frac{|\ell'|}{2}$ , then

$$\begin{aligned} \exists \delta_1 > 0: 0 < |x - x_0| < \delta_1 &\Rightarrow |g(x) - \ell'| < \frac{|\ell'|}{2} \\ &\Rightarrow ||g(x)| - |\ell'|| < \frac{|\ell'|}{2} \\ &\Rightarrow \frac{|\ell'|}{2} < |g(x)| < \frac{3|\ell'|}{2} \\ &\Rightarrow \frac{1}{|g(x)|} < \frac{2}{|\ell'|}. \end{aligned}$$

On the other hand we have:

$$\forall \varepsilon > 0; \exists \delta_2 > 0; \forall x \in V_{x_0}: 0 < |x - x_0| < \delta_2 \Rightarrow |g(x) - \ell'| < \varepsilon.$$

For  $\delta = \min\{\delta_1, \delta_2\}$ , then

$$.0 < |x - x_0| < \delta \Rightarrow \left| \frac{1}{g(x)} - \frac{1}{\ell'} \right| = \left| \frac{\ell' - g(x)}{\ell' g(x)} \right| < \frac{2|g(x) - \ell'|}{|\ell'|^2} < \frac{2\varepsilon}{|\ell'|^2} = \varepsilon'$$

**4.2.5 Indeterminate form**

We say that we are in the presence of an indeterminate form. If when  $x \rightarrow x_0$

- 1)  $f \rightarrow +\infty$  and  $g \rightarrow -\infty$  then  $f + g \rightarrow$  indeterminate form  $+\infty - \infty$ .
- 2)  $f \rightarrow \infty$  and  $g \rightarrow 0$  then  $f \cdot g \rightarrow$  indeterminate form  $\infty \cdot 0$ .
- 3)  $f \rightarrow \infty$  and  $g \rightarrow \infty$  then  $\frac{f}{g} \rightarrow$  indeterminate form  $\frac{\infty}{\infty}$ .
- 4)  $f \rightarrow 0$  and  $g \rightarrow 0$  then  $\frac{f}{g} \rightarrow$  indeterminate form  $\frac{0}{0}$ .
- 5)  $f \rightarrow 0$  and  $g \rightarrow 0$  then  $f^g \rightarrow$  indeterminate form  $0^0$ .
- 6)  $f \rightarrow \infty$  and  $g \rightarrow 0$  then  $f^g \rightarrow$  indeterminate form  $\infty^0$ .



7)  $f \rightarrow 1$  and  $g \rightarrow \infty$  then  $f^g \rightarrow$  indeterminate form  $1^\infty$ .

**Remarks**

1) The indeterminate forms  $\infty \cdot 0, \frac{\infty}{\infty}$  can be reduced to the form  $\frac{0}{0}$ . by writing  $\frac{f}{g} = \frac{1}{\frac{g}{f}}$  in (3) and  $f \cdot g =$

$\frac{g}{\frac{1}{f}}$  in (2)/

2) The indeterminate forms  $0^0, \infty^0, 1^\infty$  can be reduced to the form  $\infty \cdot 0$  by passing the logarithm.

**Examples** Calculate the limits: 1)  $\lim_{x \rightarrow -1} \frac{x^2+3x+2}{x^4+1}$ , 2)  $\lim_{x \rightarrow \infty} x \ln \frac{x+1}{x-2}$ , 3)  $\lim_{x \rightarrow \infty} \left(\frac{x+1}{x-2}\right)^x$ .

**4.2.6 Cauchy's criterion for functions:**

**Theorem 4.4**

A function  $f$  has a finite limit at  $x_0$  if and only if

$$\forall \varepsilon > 0 ; \exists \delta > 0 ; \forall x', x'' \in V_{x_0} : (0 < |x' - x_0| < \delta \text{ and } 0 < |x'' - x_0| < \delta) \Rightarrow |f(x') - f(x'')| < \varepsilon$$

**Proof**

**Necessary condition** Assume that  $\lim_{x \rightarrow x_0} f(x) = \ell$ , then

$$\forall \varepsilon > 0 ; \exists \delta > 0 ; \forall x', x'' \in V_{x_0} : (0 < |x' - x_0| < \delta \text{ and } 0 < |x'' - x_0| < \delta) \Rightarrow |f(x') - \ell| < \frac{\varepsilon}{2} \text{ and } |f(x'') - \ell| < \frac{\varepsilon}{2}$$

So

$$|f(x') - f(x'')| = |f(x') - \ell - (f(x'') - \ell)| < |f(x') - \ell| + |(f(x'') - \ell)| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

**Sufficient condition** Assume that  $\forall \varepsilon > 0 ; \exists \delta > 0 ; \forall x', x'' \in V_{x_0} :$

$$(0 < |x' - x_0| < \delta \text{ and } 0 < |x'' - x_0| < \delta) \Rightarrow |f(x') - \ell| < \frac{\varepsilon}{2} \text{ and } |f(x'') - \ell| < \frac{\varepsilon}{2}$$

Let  $(x_n)$  be a sequence of  $V_{x_0}$  elements where  $\forall n \in \mathbb{N} : x_n \neq x_0$  and  $\lim_{n \rightarrow \infty} x_n = x_0$ .

So for  $\delta > 0$ , then  $\exists N_0 \in \mathbb{N} : \forall n \in \mathbb{N} ; n > N_0 \Rightarrow |x_n - x_0| < \delta$ .

So  $\forall p, q \in \mathbb{N} : p > N_0$  and  $q > N_0 \Rightarrow 0 < |x_p - x_0| < \delta$  and  $0 < |x_q - x_0| < \delta \Rightarrow |f(x_p) - f(x_q)| < \varepsilon$ .

So  $(x_n)$  is a Cauchy sequence, and therefore convergent.

Let us now show that the limit  $\lim_{n \rightarrow \infty} f(x_n)$  is independent of the choice of sequence  $(x_n)$ .

Let  $(x_n)$  and  $(x'_n)$  where  $\lim_{n \rightarrow \infty} x'_n = \lim_{n \rightarrow \infty} x_n = x_0$ .

So  $\exists N \in \mathbb{N} ; \forall n \in \mathbb{N} : n > N \Rightarrow (0 < |x_n - x_0| < \delta \text{ and } 0 < |x'_n - x_0| < \delta) \Rightarrow |f(x_n) - f(x'_n)| < \varepsilon$ .

So

$$\lim_{n \rightarrow \infty} (f(x_n) - f(x'_n)) = 0,$$

we obtain

$$\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} f(x'_n).$$

### 4.2.7 Comparison of functions in the neighborhood of a point - Landau notation:

Let  $f$  and  $g$  be functions defined in the neighborhood  $V_{x_0}$  of the point  $x_0$ , with the possible exception of  $x_0$

#### Definition 4.8

We say that  $f$  is negligible in front of  $g$  when  $x \rightarrow x_0$ , and we write  $f = o(g)$ , if

$$\forall \varepsilon > 0; \exists \delta > 0; \forall x \in V_{x_0}: 0 < |x - x_0| < \delta \Rightarrow |f(x)| \leq \varepsilon |g(x)|.$$

#### Definition 4.9

We say that  $f$  is dominated by  $g$  when  $x \rightarrow x_0$ , and we write  $f = O(g)$ , if

$$\exists k > 0; \exists \delta > 0; \forall x \in V_{x_0}: 0 < |x - x_0| < \delta \Rightarrow |f(x)| \leq k |g(x)|.$$

The symbols  $o$  and  $O$  are called Landau symbols.

#### Corollary 4.1

If  $g$  is non-zero on  $V_{x_0} - \{x_0\}$  then:

$$f = o(g) \Leftrightarrow \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 0.$$

$$f = O(g) \Leftrightarrow \left| \frac{f(x)}{g(x)} \right| \text{ is bounded in } V_{x_0}.$$

And if  $g = 1$ , then

$$f = o(1) \Leftrightarrow \lim_{x \rightarrow x_0} f(x) = 0 \text{ and } f = O(1) \Leftrightarrow f \text{ is bounded in } V_{x_0}.$$

**Remark** We obtain a similar definition for  $x_0 = +\infty$  and  $x_0 = -\infty$ .

#### Examples

1) When  $x \rightarrow 0$  we have.

$$x^3 = o(x^2), \quad x^2 \cos \frac{1}{x} = O(x^2), \quad \left(\frac{1}{x}\right)^3 = o\left(\left(\frac{1}{x}\right)^4\right).$$

2) When  $x \rightarrow +\infty$  we have

$$x^2 = o(x^3), \quad x^2 \sin x = O(x^2), \quad \left(\frac{1}{x}\right)^4 = o\left(\left(\frac{1}{x}\right)^3\right).$$

#### Theorem 4.5

1)  $f = gh \Leftrightarrow f = o(g)$  where  $h = o(1)$ .

2)  $f = gh \Leftrightarrow f = O(g)$  where  $h = O(1)$ .

**Proof** (Let's prove 1)

**Necessary condition:** Assume that  $f = o(g)$ .

We put  $h(x) = \begin{cases} \frac{f(x)}{g(x)}, & g(x) \neq 0 \\ 0 & , g(x) = 0 \end{cases}$ .

We have  $f = o(g) \Leftrightarrow \forall \varepsilon > 0; \exists \delta > 0; \forall x \in V_{x_0}: 0 < |x - x_0| < \delta \Rightarrow |f(x)| \leq \varepsilon |g(x)|$ .

**First:** Let us prove that  $f = gh$ .

If  $g(x) = 0$  then  $0 < |x - x_0| < \delta \Rightarrow |f(x)| \leq \varepsilon |g(x)| = 0$ , we get  $f = gh$ .

If  $g(x) \neq 0$  then  $f(x) = g(x) \frac{f(x)}{g(x)}$ , we get  $f = gh$ .

**second:**

Let us show that  $h = o(1)$ , i.e  $\forall \varepsilon > 0; \exists \delta > 0; \forall x \in V_{x_0}: 0 < |x - x_0| < \delta \Rightarrow |h(x)| \leq \varepsilon$

If  $g(x) = 0$  then  $h(x)=0$ , i.e  $|h(x)| \leq \varepsilon$

If  $g(x) \neq 0$  then  $|f(x)| \leq \varepsilon |g(x)|$  and from it  $\left| \frac{f(x)}{g(x)} \right| \leq \varepsilon$  i.e  $|h(x)| \leq \varepsilon$ .

**efficient condition:**

Assume that  $f = gh$  and  $h = o(1)$  and show that  $f = o(g)$ .

We have  $(h = o(1)) \Leftrightarrow (\forall \varepsilon > 0; \exists \delta > 0; \forall x \in V_{x_0}: 0 < |x - x_0| < \delta \Rightarrow |h(x)| \leq \varepsilon)$  and

from there  $|f(x)| = |h(x)g(x)| \leq \varepsilon |g(x)|$  i.e.  $f = o(g)$ .

In the same way we prove property 2.

**Note:** The previous two properties are summarized in the following writing.

$$o(g) = g \cdot o(1) \text{ and } O(g) = g \cdot O(1)$$

**properties**

1)  $f = O(g)$  and  $h = O(g) \Rightarrow f + h = O(g)$ .

2)  $f = o(g)$  and  $h = o(g) \Rightarrow f + h = o(g)$ .

3)  $f = o(g)$  and  $h = O(1) \Rightarrow fh = o(g)$ .

4)  $f = o(g)$  and  $h = O(g) \Rightarrow f + h = O(g)$ .

5)  $f = O(g)$  and  $h = O(1) \Rightarrow fh = O(g)$ .

6)  $h = O(f)$  and  $f = o(g) \Rightarrow h = o(g)$ .

7)  $h = o(f)$  and  $f = O(g) \Rightarrow h = o(g)$ .

**Note:** The previous properties are summarized in the following writing.

1)  $O(g) + O(g) = O(g)$ .

- 2)  $o(g) + o(g) = o(g)$ .
- 3)  $o(g)O(1) = o(g)$ .
- 4)  $o(g) + O(g) = O(g)$ .
- 5)  $O(g).O(1) = O(g)$ .
- 6)  $O(o(g)) = o(g)$ .
- 7)  $o(O(g)) = o(g)$ .

#### 4.2.8 Equivalent functions:

Let  $f$  and  $g$  be functions defined in the neighborhood  $V_{x_0}$  of the point  $x_0$ , with the possible exception of  $x_0$ .

##### Definition 4.11

We say that  $f$  is equivalent to  $g$  for  $x \rightarrow x_0$  and write  $f \sim g$  if  $f - g = o(f)$  for  $x \rightarrow x_0$ .

##### Results 4.1

- 1)  $f - g = o(f) \Leftrightarrow f - g = o(g)$ .
- 2) The relation  $\sim$  is an equivalence relation on the set of functions defined in the neighborhood  $V_{x_0} - \{x_0\}$  of the point  $x_0$ .
- 3) If  $f$  and  $g$  are non-zero on  $V_{x_0} - \{x_0\}$  then:  $f \sim g \Leftrightarrow \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 1$ .

##### Theorem 4.7

Let  $f$ ,  $g$ ,  $f_1$  and  $g_1$  be functions defined in the neighborhood  $V_{x_0}$  of the point  $x_0$ , with the possible exception of  $x_0$  where  $f \sim f_1$  and  $g \sim g_1$  for  $x \rightarrow x_0$ . If

if the limit  $\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)}$  exists then the limit  $\lim_{x \rightarrow x_0} \frac{f_1(x)}{g_1(x)}$  also exists and we have:

$$\lim_{x \rightarrow x_0} \frac{f_1(x)}{g_1(x)} = \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)}$$

##### Proof

Since  $\frac{f(x)}{g(x)}$  accepts a limit when  $x \rightarrow x_0$ , there is a neighborhood  $V_{x_0}$  to the point  $x_0$ , such that  $g$  is non-zero on  $V_{x_0} - \{x_0\}$  and that  $g \sim g_1$  (that is,  $|g(x)| \leq \varepsilon |g_1(x)|$ ) then  $g_1$  is also non-zero on  $V_{x_0} - \{x_0\}$  and hence

$$\begin{cases} f \sim f_1 \\ g \sim g_1 \end{cases} \Rightarrow \begin{cases} f_1 \sim f \\ g_1 \sim g \end{cases} \Rightarrow \begin{cases} f_1 = f(1 + o(1)) \\ g_1 = g(1 + o(1)) \end{cases} \Rightarrow \frac{f_1}{g_1} = \frac{f(1 + o(1))}{g(1 + o(1))}$$

And since  $\frac{(1+o(1))}{(1+o(1))} = 1 + o(1) \rightarrow 1$ , then  $\lim_{x \rightarrow x_0} \frac{f_1(x)}{g_1(x)} = \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)}$ .

**Remark**

Note: The concept of equivalent functions is used in calculating limits, especially in removing indeterminacy.

**Examples**

1) Calculate the limit  $\lim_{x \rightarrow 0} \frac{\sqrt{4+x}-2}{\sqrt[3]{x+1}-1}$ .

For  $x \rightarrow 0$  we have  $\sqrt{4+x}-2 \sim \frac{1}{2}x$  and  $\sqrt[3]{x+1}-1 \sim \frac{1}{3}x$ , and from it

$$\lim_{x \rightarrow 0} \frac{\sqrt{4+x}-2}{\sqrt[3]{x+1}-1} = \lim_{x \rightarrow 0} \frac{\frac{1}{2}x}{\frac{1}{3}x} = \frac{3}{2}.$$

2) Calculate the limit  $\lim_{x \rightarrow +\infty} \frac{\sqrt{x^2-2x+x}}{2+xe^{\frac{1}{x}}}$ .

For  $x \rightarrow +\infty$  we have  $\sqrt{x^2-2x+x} \sim 2x$  and  $2+xe^{\frac{1}{x}} \sim x$ , and from it

$$\lim_{x \rightarrow +\infty} \frac{\sqrt{x^2-2x+x}}{2+xe^{\frac{1}{x}}} = \lim_{x \rightarrow +\infty} \frac{2x}{x} = 2.$$

**4.3 Continuous functions:**

**Definition 4.12**

Let  $f$  be a function defined on the neighborhood  $V_{x_0}$  of the point  $x_0$ . We say that  $f$  is continuous at  $x_0$  if and only if:  $\lim_{x \rightarrow x_0} f(x) = f(x_0)$ . In other words

$$(f \text{ is continuous at } x_0) \Leftrightarrow (\forall \varepsilon > 0; \exists \delta > 0; \forall x \in V_{x_0}: 0 < |x - x_0| < \delta \Rightarrow |f(x) - f(x_0)| < \varepsilon).$$

Let  $f$  be a function defined on the neighborhood  $V_{x_0}$  from the right for the point  $x_0$ , we say that  $f$  is continuous at  $x_0$  from the right if and only if:  $\lim_{x \rightarrow x_0^+} f(x) = f(x_0)$ .

Let  $f$  be a function defined on the neighborhood  $V_{x_0}$  from the left for the point  $x_0$ , we say that  $f$  is continuous at  $x_0$  from the left if and only:  $\lim_{x \rightarrow x_0^-} f(x) = f(x_0)$ .

**Result 4.2**

A function  $f$  is continuous at  $x_0$  if and only if it is continuous at  $x_0$  from the right and from the left

**Examples**

1) Let the function  $f$  defined on  $\mathbb{R}$  by  $f(x) = \begin{cases} \frac{|x^2-1|}{x-1} & \text{if } x \neq 1 \\ 2 & \text{if } x = 1 \end{cases}$ .

$\lim_{x \rightarrow 1^+} f(x) = 2 = f(1) \Rightarrow f$  is continuous at  $x_0 = 1$ , from the right.

$\lim_{x \rightarrow 1^-} f(x) = -2 \neq f(1) \Rightarrow f$  is discontinuous at  $x_0 = 1$ , from the left. So  $f$  is discontinuous at  $x_0 = 1$ .

### **Definition 4.13**

Let  $I$  be an interval of  $\mathbb{R}$ .

We say that a function  $f$  is continuous on the interval  $I$  if and only if it is continuous at every point in this interval. We denote the set of continuous functions on the interval  $I$  by  $C(I)$ .

We say that the function  $f$  is continuous uniformly over the domain  $I$  if and only if

$$\forall \varepsilon > 0; \exists \delta > 0: \forall x', x'' \in I: |x' - x''| < \delta \Rightarrow |f(x') - f(x'')| < \varepsilon.$$

It is clear from the definition that every uniformly continuous function in the interval  $I$  is continuous in this interval (the opposite is not always true).

### **4.3.1 Continuous functions in a closed interval**

#### **Theorem 4.8**

Every continuous function in a closed interval  $[a, b]$  is uniformly continuous in this interval.

#### **Proof**

We assume that  $f$  is continuous and uniformly discontinuous on  $[a, b]$  i.e.

$$\exists \varepsilon > 0; \forall \delta > 0: \exists x', x'' \in [a, b]: |x' - x''| < \delta \text{ and } |f(x') - f(x'')| \geq \varepsilon.$$

We put  $\delta = \frac{1}{n} > 0$  where  $n \in \mathbb{N}^*$  and from it:

$$\exists \varepsilon > 0; \forall n \in \mathbb{N}^*; \exists x'_n, x''_n \in [a, b]: |x'_n - x''_n| < \frac{1}{n} \text{ and } |f(x'_n) - f(x''_n)| \geq \varepsilon.$$

Since the sequence  $(x'_n)$  is bounded, according to the BOLZANO-WEIERSTRASS theorem, then a subsequence  $(x'_{n_k})$  can be extracted from it that converges towards  $\bar{x}$  from  $[a, b]$  and since

$\forall k \in \mathbb{N}: |x'_{n_k} - x''_{n_k}| < \frac{1}{n_k}$ , the partial sequence  $(x''_{n_k})$  also converges towards  $\bar{x}$ , and since  $f$  is continuous at  $\bar{x}$ , then  $\lim_{k \rightarrow \infty} (f(x'_{n_k}) - f(x''_{n_k})) = f(\bar{x}) - f(\bar{x}) = 0$ . This is a contradiction because  $\forall k \in \mathbb{N}: |f(x'_{n_k}) - f(x''_{n_k})| \geq \varepsilon$ .

#### **Theorem 4.9**

Every continuous function on the closed interval  $[a, b]$ , is bounded.

**Proof**

Assume that  $f$  continuous and unbounded on the interval  $[a, b]$ , i.e.  $\forall n \in \mathbb{N}; \exists x_n \in [a, b]: |f(x_n)| > n$ .

Since the sequence  $(x_n)$  is bounded, it is possible to extract from it a partial sequence  $(x_{n_k})$  that converges towards  $\bar{x}$  from  $[a, b]$ . Since  $f$  is continuous at  $\bar{x}$ , then  $\lim_{k \rightarrow \infty} |f(x_{n_k})| = |f(\bar{x})|$ .

This is a contradiction because  $\forall k \in \mathbb{N}: |f(x_{n_k})| > n_k \geq k$ , and hence  $\lim_{k \rightarrow \infty} |f(x_{n_k})| = +\infty$ .

**Theorem 4.10**

Any continuous function on a closed interval  $[a; b]$  reaches its upper and lower bounds at least once, that is to say there is at least  $x_1$  and  $x_2$  are from the interval  $[a; b]$  where:

$$f(x_1) = \sup_{x \in [a; b]} f(x) \text{ and } f(x_2) = \inf_{x \in [a; b]} f(x).$$

**Proof**

Let  $M = \sup_{x \in [a; b]} f(x)$ . And assume that  $\forall x \in [a; b]: f(x) \neq M$  i.e.  $\forall x \in [a; b]: f(x) < M$ .

So the function  $g$  defined on  $[a; b]$  by  $\forall x \in [a; b]: g(x) = \frac{1}{M - f(x)}$  it is continuous and strictly

positive and therefore it is bounded to this interval, i.e.:  $\exists m > 0; \forall x \in [a; b]: g(x) \leq m$  or  $\exists m > 0; \forall x \in [a; b]: f(x) \leq M - \frac{1}{m}$ . This contradicts the hypothesis  $M = \sup_{x \in [a; b]} f(x)$ .

**Theorem 4.11**

Let  $f$  be a continuous function in the interval  $[a; b]$ , if the signs of  $f(a)$  and  $f(b)$  are different, then there is at least a point  $c$  in the interval  $]a; b[$  satisfies:  $f(c) = 0$ .

**Proof**

Assume that  $f(a) < 0$  and  $f(b) > 0$ . Let the set  $E = \{x \in [a; b] / f(x) > 0\}$ , then  $E \neq \emptyset$  because  $b \in E$ . We put  $\inf E = c$  and let us prove that:  $f(c) = 0$ .

Assume that  $f(c) \neq 0$  Since  $f$  is continuous at  $c$ , there exists at least a interval of the form  $I = ]c - \alpha; c + \alpha[ \subset [a; b]$  with  $\alpha > 0$ , where  $f(x)$  and  $f(c)$  have the same sign. (See Proposition 1.3). So

if  $f(c) > 0$ , then  $\forall x \in I: f(x) > 0$  by taking  $x = c - \frac{\alpha}{2}$  we get  $f(c - \frac{\alpha}{2}) > 0$  so  $c - \frac{\alpha}{2} \in E$  and therefore  $c - \frac{\alpha}{2} \geq c = \inf E$ . and this is a contradiction.

if  $f(c) < 0$ , then  $\forall x \in I: f(x) < 0$ .

We have  $\inf E = c \implies \exists x_0 \in E: c + \alpha > x_0 \geq c \implies x_0 \in I \implies f(x_0) < 0$ . This is a contradiction because  $x_0 \in E \implies f(x_0) > 0$ . So  $f(c) = 0$ .

**Theorem 4.12**

Let  $f$  be a continuous function in the interval  $[a; b]$ . For every real number  $\lambda$  between  $f(a)$  and  $f(b)$ , there exists at least one real number  $c$  of the interval  $[a; b]$  satisfies:  $f(c) = \lambda$ .

**Proof**

**case 1:** If  $\lambda = f(a)$  it is enough to take  $c = a$ , but if  $\lambda = f(b)$  it is enough to take  $c = b$ .

**case 2:** If  $\lambda \neq f(a)$  and  $\lambda \neq f(b)$ . Then the function  $g$  defined in the interval  $[a; b]$  by

$g(x) = f(x) - \lambda$ , satisfies the conditions of Theorem 4.11, So there exists at least one real number  $c$  of the interval  $[a; b]$  where  $g(c) = 0$  and from which we get  $f(c) = \lambda$ .

**Proposition 3.2**

Let  $I$  be the interval of  $\mathbb{R}$ ,  $f$  a real function

If the function  $f$  is continuous on  $I$ , then the image of the interval  $I$  by the function  $f$  is a interval of  $\mathbb{R}$ , that is, the set  $f(I)$  is a interval.

**Proof**

Let  $y_1; y_2$  be two numbers of  $f(I)$  where  $y_1 \leq y_2$  then there are at least two numbers  $x_1; x_2$  of the interval  $I$  where  $y_1 = f(x_1)$  and  $y_2 = f(x_2)$  according to the theorem 4.12 for every number  $y$  such that  $y_1 \leq y \leq y_2$ , there exists at least an number  $x$  confined between  $x_1$  and  $x_2$  ( i.e.  $x \in I$ ), where  $y = f(x)$  and hence  $y \in f(I)$ .

**4.3.2 Extension by continuity**

**Definition 4 14**

Let  $f$  be a function defined on the domain  $I$ . With exception of the point  $x_0$  of  $I$ , we assume that  $\lim_{x \rightarrow x_0} f(x) = \ell$ . Then the function  $\tilde{f}$ , defined by  $\tilde{f}(x) = \begin{cases} f(x) & ; x \neq x_0 \\ \ell & ; x = x_0 \end{cases}$ , coincides with  $f$  on  $I - \{x_0\}$  and is continuous at  $x_0$ . The function  $\tilde{f}$  is called the extension of  $f$  with continuity at  $x_0$ .

**Example**

Let  $f$  be a function defined on  $\mathbb{R}^*$  by  $f(x) = \frac{\sin 2x}{x}$ . Since  $\lim_{x \rightarrow 0} \frac{\sin 2x}{x} = 2$ , then  $f$  can be extended by continuity at  $x_0 = 0$  to the function  $\tilde{f}$  defined by:  $\tilde{f}(x) = \begin{cases} \frac{\sin 2x}{x} & ; x \neq 0 \\ 2 & ; x = 0 \end{cases}$ .

**4.3.3 properties of monotone functions on an interval:**

**Theorem 4.13**

Let  $f: ]a, b[ \rightarrow \mathbb{R}$  be a monotonic function where  $-\infty < a < b < +\infty$ , then the limits  $\lim_{x \rightarrow a^+} f(x)$  ,  $\lim_{x \rightarrow b^-} f(x)$ , are exists ( finite or infinite ) and we have

If  $f$  increasing  $\implies -\infty \leq \inf_{x \in ]a, b[} f(x) = \lim_{x \rightarrow a^+} f(x) \leq \lim_{x \rightarrow b^-} f(x) = \sup_{x \in ]a, b[} f(x) \leq +\infty$



If  $f$  decreasing  $\Rightarrow -\infty \leq \inf_{x \in ]a, b[} f(x) = \lim_{x \rightarrow b} f(x) \leq \lim_{x \rightarrow a} f(x) = \sup_{x \in ]a, b[} f(x) \leq +\infty$

**Proof**

Assume that  $f$  increasing and  $\sup_{x \in ]a, b[} f(x) = M < +\infty$  and let us prove that:  $\lim_{x \rightarrow b} f(x) = M$ .

We have  $\sup_{x \in ]a, b[} f(x) = M \Rightarrow \forall \varepsilon > 0; \exists \alpha \in ]a, b[ : M - \varepsilon < f(\alpha) \leq M$ .

By putting  $\delta = b - \alpha > 0$ , then  $b - \delta < x < b \Rightarrow \alpha < x < b \stackrel{f \text{ increasing}}{\Rightarrow} f(\alpha) \leq f(x) \Rightarrow M - \varepsilon < f(\alpha) \leq f(x) \leq M < M + \varepsilon \Rightarrow M - \varepsilon < f(x) < M + \varepsilon$ .

So  $\forall \varepsilon > 0; \exists \delta > 0 : -\delta < x - b < 0 \Rightarrow |f(x) - M| < \varepsilon$  we get  $\lim_{x \rightarrow b} f(x) = M$ .

In the same way we prove the second case.

**Corollary 4.1**

1) Let  $f : ]a, b[ \rightarrow \mathbb{R}$  be a monotonic function then:

a) If  $f$  increasing  $\Rightarrow f(a) \leq \lim_{x \rightarrow a} f(x) \leq \lim_{x \rightarrow b} f(x) \leq f(b)$ .

b) If  $f$  decreasing  $\Rightarrow f(b) \leq \lim_{x \rightarrow b} f(x) \leq \lim_{x \rightarrow a} f(x) \leq f(a)$ .

2) Let  $I$  be an interval of  $\mathbb{R}$  bounded by  $a$  and  $b$  ( $a < b$ ), and let  $f : [a, b] \rightarrow \mathbb{R}$  be an increasing function. For each  $x_0$ , where  $a < x_0 < b$  then:

a)  $-\infty < f(x_0 - 0) \leq f(x_0) \leq f(x_0 + 0) < +\infty$ .

b) If  $a \in I \Rightarrow f(a) \leq f(a + 0) < +\infty$ .

c) If  $b \in I \Rightarrow -\infty < f(b - 0) \leq f(b)$ .

**Remark**

We obtain a corollary similar to corollary 4.1 if  $f$  is decreasing over the interval  $I$ .

**Theorem 4.14**

Let  $I$  be an interval of  $\mathbb{R}$  and let  $f : [a, b] \rightarrow \mathbb{R}$  be an monotonic function Then  $f$  is continuous on  $I$  if and only if  $f(I)$  is a interval.

**Proof**

**Necessary conditions**

According to Proposition 2.3, if  $f$  is continuous, then  $f(I)$  is an interval.

**sufficient condition**

We assume  $f$  is increasing and  $f(I)$  is an interval and prove that  $f$  is continuous on  $I$ .

Suppose the opposite and let  $x_0$  be a point of discontinuity of  $f$ . As  $f$  is increasing, then at least one of the relations  $f(x_0) < f(x_0 + 0)$ ,  $f(x_0 - 0) < f(x_0)$ . is verified (corollary 4.1).

Assume, for example, that  $f(x_0) < f(x_0 + 0)$  in this case, then for each  $x$  of  $I$ , we have

$x \leq x_0 \Rightarrow f(x) < f(x_0)$  and  $x > x_0 \Rightarrow f(x) \geq f(x_0 + 0)$  that is  $]f(x_0), f(x_0 + 0)[ \cap f(I) = \emptyset$ .

Let  $x_1 \in I$  where  $x_1 > x_0$  then  $f(x_0) \in f(I)$  and  $f(x_1) \in f(I)$  and from it  $[f(x_0), f(x_1)] \subset f(I)$  (because  $f(I)$  is an interval) and since  $f(x_1) > f(x_0 + 0)$  then  $]f(x_0), f(x_0 + 0)[ \subset [f(x_0), f(x_1)]$

i.e.  $]f(x_0), f(x_0 + 0)[ \cap f(I) \neq \emptyset$ . This is a contradiction.

#### 4.4.3 The inverse function of a strictly monotonic continuous function:

##### Theorem 4.15

Let  $I$  be the interval of  $\mathbb{R}$  and  $f: I \rightarrow \mathbb{R}$  as a function.

If  $f$  is continuous and strictly monotonic over the interval  $I$ , then  $f$  in this case is bijective of the interval  $I$  to the interval  $f(I)$ . Therefore,  $f$  accepts an inverse function that we denote by  $f^{-1}$ , which in turn is defined, continuous, and strictly monotonic over the interval  $f(I)$  and has the same direction of change of  $f$ , and we have

$$\forall x \in I; \forall y \in f(I): y = f(x) \Leftrightarrow x = f^{-1}(y) \dots (*)$$

**Remark:** Relation (\*) is used to give the expression for the function  $f^{-1}$  if possible.

If  $f$  is strictly monotonic over  $I$ , it is injective, and from the definition of the set  $f(I)$ , it is surjective, so  $f$  is bijective.

$f$  is continuous,  $f(I)$  is an interval. On the other hand, as  $f$  is strictly monotonic,  $f^{-1}$  is also monotonic. Therefore,  $f^{-1}$  is continuous according to the theorem 4.14 because  $f^{-1}(f(I)) = I$  is an interval.

##### Example

Let the function  $f$  defined on the interval  $I = [0; +\infty[$  by  $f(x) = x^2 + 3$ , then  $f$  is continuous and strictly monotonic (strictly increasing) on the interval  $I = [0; +\infty[$  where  $f(I) = [3; +\infty[$  according to theorem (4.15),  $f$  is a bijective to the interval  $[0; +\infty[$  in the interval  $[3; +\infty[$ , so it accepts an inverse function  $f^{-1}$  and we have:

$$\forall x \in [0; +\infty[; \forall y \in [3; +\infty[: y = x^2 + 3 \Leftrightarrow x^2 = y - 3$$

$$\Leftrightarrow \begin{cases} x = \sqrt{y - 3} \\ x = -\sqrt{y - 3} < 0 \text{ (مرفوض)} \end{cases}$$

So  $f^{-1}(x) = \sqrt{y - 3}$ , after replacing  $x$  with  $y$ , the final definition of the inverse function  $f^{-1}$  is as follows:

$$f^{-1}: [3; +\infty[ \rightarrow [0; +\infty[ \\ x \rightarrow \sqrt{x-3}$$

**Exercise\***

Let the function  $f$  defined on  $\mathbb{R}$  by  $f(x) = \begin{cases} x^2 - 2x + 1 & \text{si } x \leq 1 \\ \frac{-x+1}{2x-1} & \text{si } x > 1 \end{cases}$ .

- 1) Prove That  $f$  is continuous and strictly monotonic over  $\mathbb{R}$ .
- 2) Concluding that  $f$  accepts an inverse function  $f^{-1}$ , write the expression  $f^{-1}(x)$  in terms of  $x$ .

**Solution**

$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^-} f(x) = f(1) = 0 \Rightarrow$  continuous at 0  $\Rightarrow f$  continuous over  $\mathbb{R}$ .

$f$  is decreasing over  $\mathbb{R}$  and  $f(\mathbb{R}) = ]-\frac{1}{2}; +\infty[$ . So

$$f^{-1}: ]-\frac{1}{2}; +\infty[ \rightarrow \mathbb{R} \\ x \rightarrow f(x) = \begin{cases} \frac{x+1}{2x+1}, & -\frac{1}{2} < x < 0 \\ 1 - \sqrt{x}, & x \geq 0 \end{cases}$$

**4.4 Differentiable functions**

**4.4.1 Definition and basic properties**

**Definition 4.15**

Let  $f$  be a function defined on the neighborhood  $V_{x_0}$  of the point  $x_0$ . We say that the function  $f$  is differentiable at  $x_0$  if and only if  $\lim_{x \rightarrow x_0} \frac{f(x)-f(x_0)}{x-x_0} = L$ , exists. We call  $L$  the *derivative* of  $f$  at  $x_0$ , and we write.  $f'(x_0) = L$ . If  $f$  is differentiable at all  $x \in I$ , then we simply say that  $f$  is *differentiable*, and then we obtain a function  $f': I \rightarrow \mathbb{R}$  The derivative is sometimes written as  $\frac{df}{dx}$  or  $\frac{dy}{dx}$  where  $y = f(x)$ .

**Remarks**

- 1) By putting  $x - x_0 = h$ , the previous limit is written as  $\lim_{h \rightarrow 0} \frac{f(x_0+h)-f(x_0)}{h} = f'(x_0)$ .
- 2) The function  $f$  is differentiable at  $x_0$  if and only if there exists a function  $\varepsilon$  defined in the neighborhood  $V_{x_0}$  to the point  $x_0$  where

$$\forall x \in V_{x_0}: f(x) - f(x_0) = (f'(x_0) + \varepsilon(x))(x - x_0), \lim_{x \rightarrow x_0} \varepsilon(x) = 0$$

If  $\lim_{x \rightarrow x_0^+} \frac{f(x)-f(x_0)}{x-x_0} = L_d$  ( $\lim_{x \rightarrow x_0^-} \frac{f(x)-f(x_0)}{x-x_0} = L_g$ , respectively), we say that the function  $f$  is differentiable at  $x_0$  from the right (from the left, respectively) And we write  $L_d = f'(x_0 + 0)$  ( $L_g = f'(x_0 - 0)$ , respectively).

**Corollary 4.2**

A function  $f$  is differentiable at  $x_0$  if and only if  $f'(x_0 - 0)$  and  $f'(x_0 + 0)$  exist and

$$f'(x_0 + 0) = f'(x_0 - 0).$$

**Example**

Let  $f$  be a function defined in  $\mathbb{R}$  by  $f(x) = |x^2 - 1|$ , let us study the differentiability of  $f$  at  $x_0 = 1$ . We have

$$\lim_{x \rightarrow 1^+} \frac{f(x)-f(1)}{x-1} = \lim_{x \rightarrow 1^+} \frac{x^2-1}{x-1} = 2 = f'(1 + 0) \text{ and } \lim_{x \rightarrow 1^-} \frac{f(x)-f(1)}{x-1} = \lim_{x \rightarrow 1^-} \frac{-(x^2-1)}{x-1} = -2 = f'(1 - 0).$$

$f$  is differentiable at  $x_0 = 1$  from the right and from the left, but it is not differentiable at  $x_0 = 1$  because  $f'(1 + 0) \neq f'(1 - 0)$ .

**Geometric interpretation**

The derivative of the function  $f$  at  $x_0$  is the slope of the line tangent to the graph of  $f$  at the point  $M_0(x_0, f(x_0))$ . Thus, the equation of this tangent line is

$$y = f'(x_0)(x - x_0) + f(x_0).$$

The left and right derivatives are also interpreted by the half-tangents to the left and right of the point  $M_0(x_0, f(x_0))$ .

**Theorem 4.16**

If  $f$  is differentiable at  $x_0$ , then  $f$  is continuous at  $x_0$ .

**Proof**

Let  $f$  be differentiable at  $x_0$  then there is a neighborhood  $V_{x_0}$  where

$$\forall x \in V_{x_0}: f(x) - f(x_0) = (f'(x_0) + \varepsilon(x))(x - x_0) \text{ and } \lim_{x \rightarrow x_0} \varepsilon(x) = 0. \text{ So}$$

$$\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = \lim_{x \rightarrow x_0} (f'(x_0) + \varepsilon(x))(x - x_0) = 0 \text{ So } f \text{ is continuous at } x_0.$$

**4.4.2 Higher order derivative**

Let  $f$  be a function differentiable on the interval  $I$ . If  $f'$  differentiable on the interval  $I$ , then we denote its derivative by  $f''$ , it is called the second derivative. In the same way, we define the successive derivatives of the function  $f$  as follows:

$$\forall n \in \mathbb{N}: f^{(n+1)}(x) = (f^{(n)}(x))' \text{ , } f^{(0)}(x) = f(x).$$

We denote the  $n$ th-order derivative of the function  $f$  by  $\frac{d^n y}{dx^n}$  or  $y^{(n)}$ , where  $y = f(x)$ .

**Exercise** Prove that:

$$1) \forall n \in \mathbb{N} : \cos^{(n)} x = \cos\left(x + \frac{\pi}{2} n\right). \quad 2) \forall n \in \mathbb{N} : \left[\frac{1}{x}\right]^{(n)} = \frac{(-1)^n n!}{x^{n+1}}.$$

**Definition 4.16**

We say of a function  $f$  defined in interval  $I$ , that it is of class  $C^n$  if it is differentiable to order  $n$  and the derivative  $f^{(n)}$  is continuous over  $I$ . We denote the set of functions of class  $C^n$  in the interval  $I$  by  $C^n(I)$ . We have a definition:

$$C^0(I) = C(I)$$

The set of infinitely differentiable functions over the interval  $I$ , we denote  $C^\infty(I)$ .

**4.4.3 Operations on differentiable functions**

**Theorem 4.17**

Let  $f$  and  $g$  be differentiable functions on the interval  $I$ , then the functions  $f + g, \alpha f, fg, \frac{f}{g}$  ( $g \neq 0$ ) are differentiable over  $I$  and we have:

$$(f + g)' = f' + g' \quad , \quad (\alpha f)' = \alpha f'$$

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2} \quad , \quad (fg)' = f'g + fg'$$

**Proof** ( Let us prove the last case )

Let  $x_0 \in I$  we have

$$\frac{\frac{f(x)-f(x_0)}{g(x)} - \frac{f(x_0)-f(x_0)}{g(x_0)}}{x-x_0} = \frac{f(x)g(x_0) - f(x_0)g(x)}{g(x)g(x_0)(x-x_0)} = \frac{\frac{f(x)-f(x_0)}{(x-x_0)}g(x_0) - f(x_0)\frac{g(x)-g(x_0)}{(x-x_0)}}{g(x)g(x_0)}$$

When  $x \rightarrow x_0$  then  $\frac{f(x)-f(x_0)}{(x-x_0)} \rightarrow f'(x_0)$  and  $\frac{g(x)-g(x_0)}{(x-x_0)} \rightarrow g'(x_0)$  and  $f(x) \rightarrow f(x_0)$  and

$$g(x) \rightarrow g(x_0). \text{ So } \frac{\frac{f(x)-f(x_0)}{g(x)} - \frac{f(x_0)-f(x_0)}{g(x_0)}}{x-x_0} \rightarrow \frac{f'(x_0)g(x_0) - f(x_0)g'(x_0)}{(g(x_0))^2}$$

**Theorem 4.18** (Leibniz formula)

If  $f$  and  $g$  admit  $n$ th derivatives on the interval  $I$  then the function  $f \cdot g$  admits an  $n$ th derivative on the interval  $I$  and we have:

$$\forall n \in \mathbb{N}: (f \cdot g)^{(n)} = \sum_{p=0}^n C_n^p f^{(n-p)} g^{(p)}.$$

**Proof**

We use proof by induction and by noting that:  $\forall n, p \in \mathbb{N} (1 \leq p \leq n-1): C_n^p = C_{n-1}^p + C_{n-1}^{p-1}$ .

**Theorem 4.19**

Let  $f$  and  $g$  be functions where  $f$  is differentiable on the interval  $I$  and  $g$  is differentiable on the interval  $f(I)$ , then the function  $g \circ f$  is differentiable on the interval  $I$  and  $(g \circ f)' = (g' \circ f)f'$ .

**Proof**

Let  $x_0 \in I$  since  $f$  is differentiable at  $x_0$  and  $g$  is differentiable at  $y_0 = f(x_0)$ , Then

$$f(x) - f(x_0) = (f'(x_0) + \varepsilon_1(x))(x - x_0) \text{ with } \lim_{x \rightarrow x_0} \varepsilon_1(x) = 0$$

and

$$g(y) - g(y_0) = (g'(y_0) + \varepsilon_2(y))(y - y_0) \text{ with } \lim_{y \rightarrow y_0} \varepsilon_2(y) = 0.$$

For  $y = f(x)$  then  $y \rightarrow y_0$  when  $x \rightarrow x_0$  (since  $f$  is continuous at  $x_0$ ) and from there

$$g(f(x)) - g(f(x_0)) = (g'(f(x_0)) + \varepsilon_2(y))(f'(x_0) + \varepsilon_1(x))(x - x_0) \text{ and}$$

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = (g'(f(x_0)) + \varepsilon_2(y))(f'(x_0) + \varepsilon_1(x))$$

For  $x \rightarrow x_0$  then  $y \rightarrow y_0$ ,  $\varepsilon_1(x) \rightarrow 0$  and  $\varepsilon_2(y) \rightarrow 0$ . So

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} \rightarrow g'(f(x_0))f'(x_0).$$

**Example**

Let the function  $h$  defined on  $\mathbb{R}_+$  by  $h(x) = \cos(3\sqrt{x} + x^2)$ . We have  $h = g \circ f$  where  $f(x) = 3\sqrt{x} + x^2$  and  $g(x) = \cos x$  and we have  $f'(x) = \frac{3}{2\sqrt{x}} + 2x$  and  $g'(x) = -\sin x$ . So

$$\begin{aligned} h'(x) &= (g' \circ f)(x)f'(x) = -\sin(3\sqrt{x} + x^2) \left( \frac{3}{2\sqrt{x}} + 2x \right) \\ &= - \left( \frac{3}{2\sqrt{x}} + 2x \right) \sin(\sqrt{x} + x^2). \end{aligned}$$

**Theorem 4.20**

If  $f$  is strictly monotonic continuous function on the interval  $I$ , and differentiable at  $x_0$  from  $I$  where  $f'(x_0) \neq 0$ , then the inverse function  $f^{-1}$  is differentiable at  $y_0 = f(x_0)$  from  $f(I)$  And we have:

$$(f^{-1})'(y_0) = \frac{1}{f'(x_0)} = \frac{1}{f'[f^{-1}(y_0)]}.$$

**Proof**

Let  $f$  is differentiable at  $x_0$  from  $I$  where  $f'(x_0) \neq 0$ , and let  $y_0$  be a point from  $f(I)$  where

$y_0 = f(x_0)$ . For every  $y$  of  $f(I)$  there is a single real number  $x$  of  $I$  where  $y = f(x)$  and since  $f$  is continuous and strictly monotonic on  $I$ , so  $f^{-1}$  is continuous and strictly monotonic on  $f(I)$  (according to the Theorem 4.15), so  $\forall y \in f(I): y \neq y_0 \Rightarrow x \neq x_0$ . and for  $y \rightarrow y_0$ , then  $x \rightarrow x_0$ .

We put  $g = f^{-1}$  then  $y_0 = f(x_0) \Leftrightarrow x_0 = g(y_0)$  and  $y = f(x) \Leftrightarrow x = g(y)$ . So

$$\lim_{y \rightarrow y_0} \frac{g(y) - g(y_0)}{y - y_0} = \lim_{y \rightarrow y_0} \frac{x - x_0}{y - y_0} = \lim_{x \rightarrow x_0} \frac{1}{\frac{y - y_0}{x - x_0}} = \frac{1}{f'(x_0)}.$$

**Examples**

1) Let  $f: ]0; +\infty[ \rightarrow \mathbb{R}$   
 $x \rightarrow f(x) = x^n$ . The function  $f$  is continuous and strictly increasing on the domain  $I = ]0; +\infty[$ , and from it,  $f$  accepts an inverse function  $f^{-1}$  defined, continuous and strictly increasing on the interval  $f(I) = ]0; +\infty[$ , denoted by  $\sqrt[n]{\cdot}$  or  $(\cdot)^{\frac{1}{n}}$  is called the function of the  $n$ th root. Since:  $\forall x \in ]0, +\infty[: (x^n)' = nx^{n-1} \neq 0$ , Then the function  $f^{-1}$  are differentiable at every number  $y$  of the interval  $]0, +\infty[$  where  $y = x^n$  and we have:

$$(f^{-1})'(y) = \frac{1}{f'(x)} = \frac{1}{nx^{n-1}} = \frac{1}{n \left( (y^{\frac{1}{n}}) \right)^{n-1}} = \frac{1}{n} y^{\frac{1}{n}-1}. \text{ So}$$

$$\forall x \in ]0, +\infty[: \left( \sqrt[n]{x} \right)' = \left( (x)^{\frac{1}{n}} \right)' = \frac{1}{n} x^{\frac{1}{n}-1}.$$

2) Let  $h: ]-\frac{\pi}{2}; \frac{\pi}{2}[ \rightarrow \mathbb{R}$ ,  $x \rightarrow h(x) = \tan x$ . The function  $h$  is continuous and strictly increasing on the domain  $I = ]-\frac{\pi}{2}; \frac{\pi}{2}[$ , and from it,  $h$  accepts an inverse function  $h^{-1}$  defined, continuous and strictly increasing on the interval  $h(I) = \mathbb{R}$ , denoted by  $\arctan$ . Since:  $\forall x \in ]-\frac{\pi}{2}; \frac{\pi}{2}[ : h'(x) = (\tan x)' = \frac{1}{\cos^2 x} \neq 0$ , Then the function  $h^{-1}$  are differentiable at every number  $y$  of set  $\mathbb{R}$  where  $y = \tan x$  and we have:  $(h^{-1})'(y) = \frac{1}{h'(x)} = \cos^2 x = \frac{1}{1+\tan^2 x} = \frac{1}{1+y^2}$ .

So

$$\forall x \in \mathbb{R}: (\arctan x)' = \frac{1}{1+x^2}.$$

#### **Theorem 4.21**

If  $f$  has an extremum at point  $x_0$  and is differentiable at  $x_0$  then  $f'(x_0) = 0$ .

#### **Proof**

The existence of  $f'(x_0)$  entails the existence and equality of  $f'(x_0 + 0)$  and  $f'(x_0 - 0)$  and we assume that  $f(x_0)$  is a maximum, then exists a neighborhood  $V_{x_0}$  of the point  $x_0$  where

$\forall x \in V_{x_0}: f(x) \leq f(x_0)$ . So

If  $x > x_0$  then  $\frac{f(x)-f(x_0)}{x-x_0} \leq 0$  and if  $x < x_0$  then  $\frac{f(x)-f(x_0)}{x-x_0} \geq 0$ . So

$$\lim_{x \rightarrow x_0} \frac{f(x)-f(x_0)}{x-x_0} = f'(x_0 - 0) = f'(x_0) \geq 0 \text{ and}$$

$$\lim_{x \rightarrow x_0} \frac{f(x)-f(x_0)}{x-x_0} = f'(x_0 + 0) = f'(x_0) \leq 0.$$

We obtain  $f'(x_0) = 0$

#### **4.4.4 The theorems of Lagrange and Cauchy on finite increments**

##### **Proposition 3.3** (Rolle's Theorem)

If a function  $f [a, b] \rightarrow \mathbb{R}$  is continuous on a closed interval  $[a, b]$  and differentiable on the open interval  $]a, b[$  and  $f(a) = f(b)$ , then there exists a point  $c \in [a, b]$  such that  $f'(c) = 0$ .

#### **Proof**

Since the function  $f$  is continuous on  $[a, b]$ , there exist points  $x_m, x_M \in [a, b]$  where they take their minimum and maximum values respectively. If  $f(x_m) = f(x_M)$ , then the function is constant on  $[a, b]$ ; and since in that case  $\forall x \in ]a; b[: f'(x) = 0$ . If  $f(x_m) < f(x_M)$ , then, since  $f(a) = f(b)$ , one of the points  $x_m$  and  $x_M$  must lie in the open interval  $]a, b[$ . We denote it by  $c$ . According theorem 4.21 we obtain  $f'(c) = 0$ .

##### **Theorem 4 22** (Lagrange's finite-increment theorem)

If a function  $f [a, b] \rightarrow \mathbb{R}$  is continuous on a closed interval  $[a, b]$  and differentiable on the open interval  $]a, b[$ , then there exists a point  $c \in [a, b]$  such that  $f(b) - f(a) = f'(c)(b - a)$ .

#### **Proof**

It is sufficient to check that the function  $g$ , defined in the domain  $[a, b]$  by  $g(x) = f(x) - \frac{f(b)-f(a)}{b-a}x$ , satisfies the conditions of Proposition 3.3. Then there is at least a number  $c$  of the interval  $]a, b[$  that satisfies  $g'(c) = 0$  and we obtain  $f'(c) = \frac{f(b)-f(a)}{b-a}$ .

**Remark**

This theorem is used in approximate calculations and in proving many inequalities.

**Example**

Using the finite increment theorem, prove that:  $\forall x \geq 0: \ln(x + 1) \leq x$ .

Applying the theorem of finite increments to the interval  $[0; x]$  where  $x \geq 0$ , we get

$$\forall x \geq 0 : \ln(x + 1) - \ln 1 = f'(c)(x - 0) ; \quad 0 < c < x.$$

So

$$\ln(x + 1) = f'(c)x = \frac{1}{1 + c} \cdot x \quad ; \quad 0 < c < x.$$

We have

$$c > 0 \Rightarrow \frac{1}{1 + c} < 1 \Rightarrow \frac{1}{1 + c}x \leq x.$$

We obtain

$$\forall x \geq 0 : \ln(x + 1) \leq x.$$

**Theorem 4 23** (Cauchy's finite-increment theorem)

If a functions  $f, g [a, b] \rightarrow \mathbb{R}$  are continuous on a closed interval  $[a, b]$  and differentiable on the open interval  $]a, b[$ , and  $g'$  is non-zero in the interval  $]a, b[$  then there exists a point  $c \in ]a, b[$  such that  $\frac{f(b)-f(a)}{g(b)-g(a)} = \frac{f'(c)}{g'(c)}$ .

**Proof**

We have  $(\forall x \in ]a ; b[: g'(x) \neq 0) \Rightarrow (g(b) \neq g(a))$  so it is sufficient to check that the function  $\varphi$ , defined in the domain  $[a, b]$  by  $\varphi(x) = f(x) - \frac{f(b)-f(a)}{g(b)-g(a)}g(x)$ , satisfies the conditions of Proposition 3.3. Then there is at least a number  $c$  of the interval  $]a, b[$  that satisfies  $\varphi'(c) = 0$  and we obtain  $\frac{f'(c)}{g'(c)} = \frac{f(b)-f(a)}{b-a}$ .

**Theorem 4 24** (Hospital Rule)

If a functions  $f, g$  are continuous on a neighborhood  $V_a$  of the point  $a$  and differentiable on  $V - \{a\}$  then: If the  $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$  exists, then the  $\lim_{x \rightarrow a} \frac{f(x)-f(a)}{g(x)-g(a)}$  also and  $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = \lim_{x \rightarrow a} \frac{f(x)-f(a)}{g(x)-g(a)}$ . If in particular,  $f(a) = g(a) = 0$  we have the equality  $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = \lim_{x \rightarrow a} \frac{f(x)}{g(x)}$ .

**Proof**



Assume that  $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = \ell$ .

For  $x > a$  we apply Theorem 4.24 to the interval  $[a, x]$  and we get:

$$\frac{f(x) - f(a)}{g(x) - g(a)} = \frac{f'(c)}{g'(c)} \text{ where } c \in ]a, x[.$$

$$\text{So } x \xrightarrow{>} a \Rightarrow c \xrightarrow{>} a \Rightarrow \frac{f'(c)}{g'(c)} \rightarrow \ell \Rightarrow \frac{f(x) - f(a)}{g(x) - g(a)} \rightarrow \ell$$

For  $x < a$  we apply Theorem 4.24 to the interval  $[x, a]$  and we get:

$$\text{So } x \xrightarrow{<} a \Rightarrow c \xrightarrow{<} a \Rightarrow \frac{f'(c)}{g'(c)} \rightarrow \ell \Rightarrow \frac{f(x) - f(a)}{g(x) - g(a)} \rightarrow \ell.$$

$$\text{We obtain } \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \ell.$$

### Remarks

- 1) The previous result remains true if  $f$  and  $g$  are undefined at  $a$  but accept two finite limits.
- 2) Theorem 4.24 can be applied several times in a row.
- 3) Theorem 4.24 can be applied in the following cases:

a)  $\lim_{x \rightarrow \infty} f(x) = 0$  and  $\lim_{x \rightarrow \infty} g(x) = 0$ .

b)  $\lim_{x \rightarrow a} f(x) = \infty$  and  $\lim_{x \rightarrow a} g(x) = \infty$ .

c)  $\lim_{x \rightarrow \infty} f(x) = \infty$  and  $\lim_{x \rightarrow \infty} g(x) = \infty$ .

### Examples

1)  $\lim_{x \rightarrow 1} \frac{\sqrt{x+3}-2}{x-1}$  ( I.F  $\frac{0}{0}$  ).

$$\lim_{x \rightarrow 1} \frac{\sqrt{x+3}-2}{x-1} = \lim_{x \rightarrow 1} \frac{1}{2\sqrt{x+3}} = \frac{1}{4}.$$

2)  $\lim_{x \rightarrow 0} \frac{e^x - x - 1}{x^2}$  ( I.F  $\frac{0}{0}$  ).

$$\lim_{x \rightarrow 0} \frac{e^x - x - 1}{x^2} = \lim_{x \rightarrow 0} \frac{e^x - 1}{2x} = \lim_{x \rightarrow 0} \frac{e^x}{2} = \frac{1}{2}.$$

3)  $\lim_{x \rightarrow +\infty} \frac{e^x + x^2}{x^3 - x + 1}$  ( I.F  $\frac{\infty}{\infty}$  ).

$$\lim_{x \rightarrow +\infty} \frac{e^x + x^2}{x^3 - x + 1} = \lim_{x \rightarrow +\infty} \frac{e^x + 2x}{3x^2 - 1} = \lim_{x \rightarrow +\infty} e^x \frac{e^x + 1}{6x} = \lim_{x \rightarrow +\infty} \frac{e^x}{6} = +\infty.$$

4)  $\lim_{x \rightarrow +\infty} \frac{2x^2}{x+3} \ln \frac{x-1}{x+2}$  ( I.F  $\infty \cdot 0$  )

$$\lim_{x \rightarrow +\infty} \frac{2x^2}{x+3} \ln \frac{x-1}{x+2} = \lim_{x \rightarrow +\infty} \frac{2x}{x+3} \lim_{x \rightarrow +\infty} \frac{\ln \frac{x-1}{x+2}}{\frac{1}{x}}.$$

Calculate  $\lim_{x \rightarrow +\infty} \frac{\ln \frac{x-1}{x+2}}{\frac{1}{x}}$  (I.F  $\frac{0}{0}$ ).

$$\lim_{x \rightarrow +\infty} \frac{\ln \frac{x-1}{x+2}}{\frac{1}{x}} = \lim_{x \rightarrow +\infty} \frac{\left(\ln \frac{x-1}{x+2}\right)'}{\left(\frac{1}{x}\right)'} = \lim_{x \rightarrow +\infty} \frac{\frac{3}{(x+2)(x-1)}}{-\frac{1}{x^2}} = -3$$

So  $\lim_{x \rightarrow +\infty} \frac{2x^2}{x+3} \ln \frac{x-1}{x+2} = 2 \times (-3) = -6$ .

## Chapter five: Elementary functions

### 5.1 Inverse Trigonometric functions

#### 5.1.1 Arcsine Function

##### *Definition 5.1*

The function  $f$  defined in the interval  $I = \left[-\frac{\pi}{2}; \frac{\pi}{2}\right]$  by  $f(x) = \sin x$ , is continuous and strictly increasing in the interval  $I$ , it accepts an inverse function  $f^{-1}$  that is defined, continuous and strictly increasing on the interval  $f(I) = [-1; 1]$ . We denote the function  $f^{-1}$  by "arcsin" or " $\sin^{-1}$ ".

We have  $\forall x \in \left[-\frac{\pi}{2}; \frac{\pi}{2}\right]; \forall y \in [-1; 1] : y = \sin x \Leftrightarrow x = \arcsin y$ .

##### *Derived function*

We have  $\forall x \in \left]-\frac{\pi}{2}; \frac{\pi}{2}\right[ : (\sin x)' = \cos x \neq 0$  ( $\cos x > 0$ )

According to the theorem 4.20 then, the function arcsin is differentiable at every number  $y$  of the field  $] -1; 1[$  where  $y = \sin x$  and we have:

$$(\arcsin y)' = \frac{1}{(\sin x)'} = \frac{1}{\cos x} = \frac{1}{\sqrt{1 - \sin^2 x}} = \frac{1}{\sqrt{1 - y^2}}$$

So

$$\forall x \in ] -1; 1[ : (\arcsin x)' = \frac{1}{\sqrt{1 - x^2}}$$

#### 5.1.2 Arccosine Function

##### *Definition 5.2*

The function  $g$  defined in the interval  $I = [0; \pi]$  by  $g(x) = \cos x$ , is continuous and strictly decreasing in the interval  $I$ , it accepts an inverse function  $g^{-1}$  that is defined, continuous and strictly decreasing on the interval  $f(I) = [-1; 1]$ . We denote the function  $g^{-1}$  by "arccos" or " $\cos^{-1}$ ".

We have  $\forall x \in [0; \pi]; \forall y \in [-1; 1] : y = \cos x \Leftrightarrow x = \arccos y$ .

##### *Derived function*

We have  $\forall x \in ]0; \pi[: (\cos x)' = -\sin x \neq 0$  ( $\sin x > 0$ ).

Then the function arccos is differentiable at every number  $y$  of the field  $] -1; 1[$  where  $y = \cos x$  and we have:

$$(\arccos y)' = \frac{1}{(\cos x)'} = -\frac{1}{\sin x} = -\frac{1}{\sqrt{1 - \cos^2 x}} = -\frac{1}{\sqrt{1 - y^2}}$$

So

$$\forall x \in ] -1; 1[ : (\arccos x)' = -\frac{1}{\sqrt{1 - x^2}}$$

### 5.1.3 Arctangent Function

#### Definition 5.3

The function  $h$  defined in the interval  $I = ]-\frac{\pi}{2}; \frac{\pi}{2}[$  by  $h(x) = \tan x$ , is continuous and strictly increasing in the interval  $I$ , it accepts an inverse function  $h^{-1}$  that is defined, continuous and strictly increasing on the interval  $h(I) = \mathbb{R}$ . We denote the function  $h^{-1}$  by "arctan" or " $\tan^{-1}$ ".

We have  $\forall x \in ]-\frac{\pi}{2}; \frac{\pi}{2}[; \forall y \in \mathbb{R} : y = \tan x \Leftrightarrow x = \arctan y$ .

#### Derived function

We have  $\forall x \in ]-\frac{\pi}{2}; \frac{\pi}{2}[ : (\tan x)' = \frac{1}{\cos^2 x} \neq 0$

Then, the function arctan is differentiable at every number  $y$  of  $\mathbb{R}$  where  $y = \tan x$  and we have:

$$(\arctan y)' = \frac{1}{(\tan x)'} = \cos^2 x = \frac{1}{1 + \tan^2 x} = \frac{1}{1 + y^2}$$

So

$$\forall x \in \mathbb{R} : (\arctan x)' = \frac{1}{1 + x^2}$$

### 5.1.4 Arccotangent Function

#### Definition 5.4

The function  $k$  defined in the interval  $I = ]0; \pi[$  by  $k(x) = \cotan x$ , is continuous and strictly decreasing in the interval  $I$ , it accepts an inverse function  $k^{-1}$  that is defined, continuous and strictly decreasing on the interval  $k(I) = \mathbb{R}$ . We denote the function  $k^{-1}$  by "arccotan" or " $\cotan^{-1}$ ".

We have  $\forall x \in ]0; \pi[; \forall y \in \mathbb{R} : y = \cotan x \Leftrightarrow x = \text{arccotan } y$ .

#### Derived function

We have  $\forall x \in ]0; \pi[ : (\cotan x)' = -\frac{1}{\sin^2 x} \neq 0$

Then, the function arccotan is differentiable at every number  $y$  of  $\mathbb{R}$  where  $y = \cotan x$  and we have:

$$(\operatorname{arccotan} y)' = \frac{1}{(\cotan x)'} = -\sin^2 x = -\frac{1}{1 + \cotan^2 x} = -\frac{1}{1 + y^2}.$$

So

$$\forall x \in \mathbb{R} : (\operatorname{arccotan} x)' = -\frac{1}{1 + x^2}.$$

### **Properties**

1)  $\forall x \in [-1; 1] : \arcsin x + \arccos x = \frac{\pi}{2}.$

2)  $\forall x \in [-1; 1] : \sin(\arccos x) = \sqrt{1 - x^2}.$

3)  $\forall x \in [-1; 1] : \cos(\arcsin x) = \sqrt{1 - x^2}.$

4)  $\forall x \in \mathbb{R} : \arctan x + \operatorname{arccotan} x = \frac{\pi}{2}.$

5)  $\forall x > 0 : \arctan x + \arctan \frac{1}{x} = \frac{\pi}{2}.$

6)  $\forall x < 0 : \arctan x + \arctan \frac{1}{x} = -\frac{\pi}{2}.$

### **Proof**

1) We put  $\forall x \in [-1; 1] : f(x) = \arcsin x + \arccos x.$

We have  $\forall x \in ]-1; 1[ : f'(x) = \frac{1}{\sqrt{1-x^2}} - \frac{1}{\sqrt{1-x^2}} = 0.$  So the function  $f$  is constant in the interval  $[-1; 1].$  So  $\forall x \in [-1; 1] : f(x) = f(0) = \frac{\pi}{2}.$

2) We have  $\forall x \in [-1; 1] : \arcsin x \in \left[-\frac{\pi}{2}; \frac{\pi}{2}\right] \Rightarrow \cos(\arcsin x) \geq 0.$  So

$$\forall x \in [-1; 1] : \cos(\arcsin x) = \sqrt{1 - (\sin(\arcsin x))^2} = \sqrt{1 - x^2}.$$

6) We put  $\forall x < 0 : f(x) = \arctan x + \arctan \frac{1}{x}.$  We have

$\forall x < 0 : f'(x) = \frac{1}{1+x^2} - \frac{1}{x^2} \frac{1}{1+(\frac{1}{x})^2} = 0.$  So the function  $f$  is constant in the interval  $]-\infty; 0[.$  So

$$\forall x \in ]-\infty; 0[ : f(x) = f(-1) = -\frac{\pi}{4} - \frac{\pi}{4} = -\frac{\pi}{2}.$$

**Remark:** The properties of inverse trigonometric functions are deduced from the properties of trigonometric functions. For example, property 1 is deduced from the property:  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$ , which we will explain later.

We have  $\frac{\pi}{2} - \alpha \in \left[-\frac{\pi}{2}; \frac{\pi}{2}\right] \Leftrightarrow \alpha \in [0, \pi]$ . Bu putting  $\cos \alpha = x$  we get  $\alpha \in [0, \pi] \Leftrightarrow x \in [-1; 1]$

and  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha \Leftrightarrow \sin\left(\frac{\pi}{2} - \alpha\right) = x \Leftrightarrow \frac{\pi}{2} - \alpha = \arcsin x$

$$\Leftrightarrow \frac{\pi}{2} - \arccos x = \arcsin x$$

$$\Leftrightarrow \frac{\pi}{2} = \arccos x + \arcsin x$$

## 5.2 Hyperbolic functions and their inverses

### 5.2.1 Hyperbolic functions

**Definition 5.5** The hyperbolic sine function, which we denote by “sh,” is defined as  $\forall x \in \mathbb{R}$

$$\mathbb{R}: \operatorname{sh} x = \frac{e^x - e^{-x}}{2}.$$

**Definition 5.6** The hyperbolic cosine function, which we denote by “ch,” is defined as  $\forall x \in \mathbb{R}$

$$\mathbb{R}: \operatorname{ch} x = \frac{e^x + e^{-x}}{2}.$$

**Definition 5.7** The hyperbolic tangent function, which we denote by “th,” is defined as

$$\forall x \in \mathbb{R}: \operatorname{th} x = \frac{\operatorname{sh} x}{\operatorname{ch} x} = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$

**Definition 5.8** The hyperbolic cotangent function, which we denote by “th,” is defined as

$$\forall x \in \mathbb{R}^*: \operatorname{coth} x = \frac{\operatorname{ch} x}{\operatorname{sh} x} = \frac{e^x + e^{-x}}{e^x - e^{-x}}.$$

### Properties

For all  $x, y \in \mathbb{R}$  we have:

$$1) \operatorname{sh}(-x) = -\operatorname{sh} x \quad \operatorname{ch}(-x) = \operatorname{ch} x.$$

$$2) 1 - \operatorname{th}^2 x = \frac{1}{\operatorname{ch}^2 x}, \operatorname{ch}^2 x - \operatorname{sh}^2 x = 1.$$

$$3) \operatorname{ch}(x + y) = \operatorname{ch} x \operatorname{ch} y + \operatorname{sh} x \operatorname{sh} y.$$

$$4) \operatorname{sh}(x + y) = \operatorname{ch} x \operatorname{sh} y + \operatorname{sh} x \operatorname{ch} y.$$

$$5) \operatorname{th}(x + y) = \frac{\operatorname{th} x + \operatorname{th} y}{1 + \operatorname{th} x \operatorname{th} y}.$$

$$6) (\operatorname{sh} x)' = \operatorname{ch} x, (\operatorname{ch} x)' = \operatorname{sh} x, (\operatorname{th} x)' = \frac{1}{\operatorname{ch}^2 x}, (\operatorname{coth} x)' = -\frac{1}{\operatorname{sh}^2 x}.$$

## 5.2.2 Inverses Hyperbolic functions

### Definition 5.9

The function  $f$  defined in the interval  $I = [0; +\infty[$  by  $f(x) = \operatorname{ch} x$ , is continuous and strictly increasing in the interval  $I$ , it accepts an inverse function  $f^{-1}$  that is defined, continuous and strictly increasing on the interval  $f(I) = [1; +\infty[$ . We denote the function  $f^{-1}$  by "arg ch" or " $\operatorname{ch}^{-1}$ ".

We have  $\forall x > 0; \forall y > 1 : y = \operatorname{ch} x \Leftrightarrow \operatorname{ch} x = \frac{e^x + e^{-x}}{2} \Leftrightarrow e^{2x} - 2ye^x + 1 = 0$ .

$$\Leftrightarrow \begin{cases} x = \ln \left( y + \sqrt{y^2 - 1} \right) \\ x = \ln \left( y - \sqrt{y^2 - 1} \right) \end{cases}$$

$$\Leftrightarrow x = \ln \left( y - \sqrt{y^2 - 1} \right) \quad \left( \text{because } \ln \left( y - \sqrt{y^2 - 1} \right) \leq 0 \right).$$

So  $\forall x \geq 1 : \operatorname{arg ch} x = \ln(x + \sqrt{x^2 - 1})$ .

**Derived function:**  $\forall x \in ]1; +\infty[ : (\operatorname{arg ch} x)' = \frac{1}{\sqrt{x^2 - 1}}$ .

### Definition 5.10

The function  $g$  defined in the interval  $I = \mathbb{R}$  by  $g(x) = \operatorname{sh} x$ , is continuous and strictly increasing in the interval  $I$ , it accepts an inverse function  $g^{-1}$  that is defined, continuous and strictly increasing on the interval  $f(I) = \mathbb{R}$ . We denote the function  $g^{-1}$  by "arg sh" or " $\operatorname{sh}^{-1}$ ".

We have  $\forall x \in \mathbb{R} : \operatorname{arg sh} x = \ln(x + \sqrt{x^2 + 1})$ .

**Derived function:**  $\forall x \in \mathbb{R} : (\operatorname{arg sh} x)' = \frac{1}{\sqrt{x^2 + 1}}$ .

### Definition 5.11

The function  $h$  defined in the interval  $I = \mathbb{R}$  by  $h(x) = \operatorname{th} x$ , is continuous and strictly increasing in the interval  $I$ , it accepts an inverse function  $h^{-1}$  that is defined, continuous and strictly increasing on the interval  $h(I) = ]-1; 1[$ . We denote the function  $h^{-1}$  by "arctan" or " $\operatorname{tan}^{-1}$ ".

We have  $\forall x \in ]-1; 1[ : \operatorname{arg th} x = \frac{1}{2} \ln \frac{1+x}{1-x}$ .

**Derived function:**  $\forall x \in ]-1; 1[ : (\operatorname{arg th} x)' = \frac{1}{1-x^2}$ .