

## Chapter 2

|                                                             |          |
|-------------------------------------------------------------|----------|
| <b>2. First order ordinary differential equations</b> ..... | <b>2</b> |
| 2.1 Generalities .....                                      | 2        |
| 2.2 Separable Equations .....                               | 3        |
| 2.3 Homogeneous Equations .....                             | 4        |
| 2.4 First-Order Linear differential Equations .....         | 5        |
| 2.5 Bernoulli Equations .....                               | 7        |

# First order ordinary differential equations

## 2.1 Generalities

### Definition 2.1

We call a first-order differential equation, each equation of the form

$$F(x, y, y') = 0 \dots \dots \dots (*).$$

where  $y: I \rightarrow \mathbb{R}$  is a function of the variable  $x$  defined in the interval  $I$  and  $y'$  is the first derivative of the function  $y$  and  $F: (x, y, z) \rightarrow F(x, y, z)$  is a function of the variables  $x, y, z$ .

### Definition 2.2

We call the solution of the differential equation  $(*)$ , in the interval  $I$ , each function  $\phi$  is differentiable in the interval  $I$  and satisfies the following.

$$F(x, \phi(x), \phi'(x)) = 0.$$

### Definition 2.3

The general solution of the differential equation in the interval  $I$  it represents the set of all solutions, i.e., the set of all functions which satisfy the equation in the interval  $I$ .

### Example 2.1

$$y' - \frac{2}{x}y - \frac{1}{2}x\sqrt{x} = 0 \dots \dots (1)$$

(1) is a first-order differential equation.

The function  $\phi: ]0, +\infty[ \rightarrow \mathbb{R}$  where  $\phi(x) = x^2\sqrt{x}$  is a solution to equation (1) because  $\phi$  is differentiable over the interval  $]0, +\infty[$  and achieves  $\phi'(x) - \frac{2}{x}\phi(x) - \frac{1}{2}x\sqrt{x} = 0$ .

Indeed

$$\phi(x) = x^2\sqrt{x}; \quad \phi'(x) = \frac{5}{2}x\sqrt{x}.$$

So

$$\phi'(x) - \frac{2}{x}\phi(x) - \frac{1}{2}x\sqrt{x} = \frac{5}{2}x\sqrt{x} - \frac{2}{x}x^2\sqrt{x} - \frac{1}{2}x\sqrt{x} = \frac{5}{2}x\sqrt{x} - \frac{2}{x}x\sqrt{x} - \frac{1}{2}x\sqrt{x} = 0.$$

There are different types of first-order differential equations, but we will only mention four of them.

## 2.2 Separable Equations.

### Definition 2.4

The first order ODE  $y' = f(x, y)$  is said to be separable if  $f(x, y)$  can be expressed as a product of a function of  $x$  times a function of  $y$ . That is, the equation is separable if the function  $f$  has the form:  $f(x, y) = g(x)h(y)$ , where  $g$  and  $h$  are continuous functions on some interval  $I$ .

The solution method is based on writing the equation in the form

$$\frac{1}{h(y)} dy = g(x) dx.$$

Of course, in dividing the equation by  $h(y)$  we have to assume that  $h(y) \neq 0$ .

### Example 2.2

The differential equation  $y' = -\frac{x}{y}$  is separable since

$$f(x, y) = -\frac{x}{y} = (-x) \left(\frac{1}{y}\right).$$

Writing the equation in the form

$$y dy = x dx$$

and integrating

$$\int y dy = \int x dx$$

we get

$$x^2 + y^2 = C.$$

### Example 2.3

Show that the differential equation

$$y' = \frac{yx - y}{y + 1},$$

is separable. Then

1. Find the general solution and any singular solutions.
2. Find a solution which satisfies the initial condition  $y(2) = 1$ .

We have  $f(x, y) = (x - 1) \frac{y}{y+1}$ .

For  $y \neq 0$ , writing the equation in the form:

$$\left(\frac{1}{y} + 1\right) dy = (x - 1) dx$$

Integrating with respect to  $x$ , we get

$$\int \left(\frac{1}{y} + 1\right) dy = \int (x - 1) dx$$

and

$$\ln|y| + y = \frac{1}{2}x^2 - x + C.$$

Is the general solution. Again we have  $y$  defined implicitly as a function of  $x$ . Note that  $y = 0$  is a solution of the differential equation (verify this), but this function is not included in the general solution

To find a solution that satisfies the initial condition, set  $x = 2, y = 1$  in the general solution:

$$\ln|1| + 1 = \frac{1}{2}(2)^2 - 2 + C \text{ which implies } C = 1.$$

A particular solution that satisfies the initial condition is:  $\ln|y| + y = \frac{1}{2}x^2 - x + 1$ .

## 2.3 Homogeneous Equations

### Definition 2.4

An ordinary differential equation  $y' = f(x, y)$ , is said to be a homogeneous differential equation if the following condition is satisfied  $f(\lambda x, \lambda y) = f(x, y)$ , for any  $\lambda \in \mathbb{R}$ .

Set  $y = vx$ ; thus the general form of first order **ODE** becomes

$$y' = \frac{d(vx)}{dx} = v + xv' = f(x, vx)$$

On obtain

$$v' = \frac{f(x, vx) - v}{x} \dots \dots \dots (*)$$

We can use variable separation to solve the equation (\*).

### Example 2.4

Find the solution of the following equation

$$y' = \frac{y^2 + 2xy}{x^2}.$$

Set

$$f(x, y) = \frac{y^2 + 2xy}{x^2}$$

Clearly,

$$f(\lambda x, \lambda y) = \frac{(\lambda y)^2 + 2\lambda x \lambda y}{(\lambda x)^2} = f(x, y).$$

Therefore, this equation is homogenous.

Now to find the solution, we set  $y = vx$ ; and the equation can be written as follows

$$v' = \frac{f(x, vx) - v}{x} = \frac{\frac{v^2x^2 + 2xvx}{x^2} - v}{x} = \frac{v^2 + v}{x}.$$

By inspection,  $v = -1$  and  $v = 0$  are solutions ( i.e.  $y = -x$  and  $y = 0$  ).

For  $v \neq -1$  and  $v \neq 0$  we have

$$\frac{dx}{x} = \frac{dv}{v^2 + v},$$

if you integrate the two sides, we get

$$\ln|x| + c = \int \frac{1}{v^2 + v} dv = \int \frac{1}{v} - \frac{1}{v + 1} dv = \ln|v| - \ln|v + 1|$$

so

$$\ln|x| + c = \ln \left| \frac{v}{v + 1} \right|$$

we get

$$\frac{v}{v + 1} = kx$$

so

$$v = \frac{kx}{1 - kx}$$

There fore,

$$y = \frac{kx^2}{1 - kx}$$

So the general solution of the original differential equation, is

$$y = \frac{kx^2}{1 - kx} \text{ or } y = -x.$$

### Example 2.5

Find the solution of the following equation

$$y' = \frac{1}{x^2}y^2 + \frac{1}{x}y + 1.$$

Set

$$f(x, y) = \frac{1}{x^2}y^2 + \frac{1}{x}y + 1.$$

Clearly,

$$f(x, y) = \frac{1}{(\lambda x)^2}(\lambda y)^2 + \frac{1}{(\lambda x)}(\lambda y) + 1 = f(x, y).$$

Now to find the solution, we set  $y = vx$ ; and the equation can be written as follows

$$v' = \frac{f(x, vx) - v}{x} = \frac{\frac{1}{x^2}(xv)^2 + \frac{1}{x}(xv) + 1 - v}{x} = \frac{v^2 + 1}{x}.$$

So

$$\frac{dv}{v^2 + 1} = \frac{dx}{x}$$

if you integrate the two sides, we get

$$\int \frac{1}{v^2 + 1} dv = \int \frac{1}{x} dx.$$

So

$$\arctan v = \ln x + C.$$

Or

$$v = \tan(\ln x + C).$$

So the general solution of the original differential equation, is

$$y = x \tan(\ln x + C).$$

## 2.4 First-Order Linear differential Equations

### Definition 2.5

A first-order linear differential equation is one that can be written in the form

$$y' - a(x)y = b(x) \dots \dots \dots (1)$$

where  $a$  and  $b$  are continuous functions of  $x$ . Equation (1) is the linear equation's standard form.

### Solving Linear Equations

We solve the equation

$$y' + a(x)y = b(x).$$

By multiplying both sides by a *positive* function  $v(x)$  that transforms the left-hand side into the derivative of the product  $v(x)y$ . We will show how to find the function  $v$  later, but first we want to show how, once found, it provides the solution we seek.

Multiplying by  $v(x)$  gives:

So  $v(x)y' + a(x)v(x)y = v(x)b(x)$  ( $v(x)$  is chosen to make  $vy' + avy = (vy)'$ ).

$$(v(x)y)' = \frac{d}{dx}(v(x)y) = v(x)b(x)$$

$$v(x)y = \int v(x)b(x) dx.$$

So the solution would be

$$y = \frac{1}{v(x)} \int v(x)b(x) dx \dots \dots \dots (2)$$

Equation (2) expresses the solution of Equation (1) in terms of the function  $v(x)$  and  $b(x)$ . We call  $v(x)$  an **integrating factor** for Equation (1).

To find such an  $v$ , we have:

$$\frac{d}{dx}(vy) = v \frac{dy}{dx} + avy$$

$$v \frac{dy}{dx} + y \frac{dv}{dx} = v \frac{dy}{dx} + avy$$

$$\frac{dv}{dx} = av.$$

This is a separable differential equation for  $v$ , which we solve as follows:

$$\int \frac{dv}{v} = \int a dx$$

$$\ln v = \int a dx$$

$$v(x) = e^{\int a(x)dx}.$$

To solve the linear differential equation  $y' + a(x)y = b(x)$ , multiply both sides by the **integrating factor**  $v(x) = e^{\int a(x)dx}$  and integrate both sides.

When you integrate the left-hand side product in this procedure, you always obtain the product  $v(x)y$  of the integrating factor and solution function  $y$  this is after defining  $v$ .

**Example 2.6**

Solve the differential equation  $\frac{dy}{dx} + 3x^2y = 6x^2$ .

**Solution**

The given equation is linear since it has the form of Equation (1) with  $a(x) = 3x^2$  and  $b(x) = 6x^2$ . An integrating factor is  $v(x) = e^{\int a(x)dx} = e^{\int 3x^2dx} = e^{x^3}$ . Multiplying both sides of the differential equation by  $e^{x^3}$ , we get

$$e^{x^3} \frac{dy}{dx} + 3x^2 e^{x^3} y = 6x^2 e^{x^3}$$

or

$$\frac{d}{dx}(e^{x^3} y) = 6x^2 e^{x^3}$$

Integrating both sides, we have

$$e^{x^3} y = \int 6x^2 e^{x^3} dx = 2e^{x^3} + C$$

$$y = 2 + C e^{-x^3}.$$

**Example 2.7**

Find the solution of the initial-value problem  $x^2y' + xy = 1$ ;  $x > 0$ ;  $y(1) = 2$

**Solution**

We must first divide both sides by the coefficient of  $y'$  to put the differential equation into standard form:  $y' + \frac{1}{x}y = \frac{1}{x^2}$ ;  $x > 0 \dots \dots \dots (4)$

The integrating factor is

$$v(x) = e^{\int a(x)dx} = e^{\int \frac{1}{x} dx} = e^{\ln x} = x$$

Multiplication of Equation ( 4 ) by  $x$  gives

$$xy' + y = \frac{1}{x} \text{ or } (xy)' = \frac{1}{x}.$$

Then

$$xy = \int \frac{1}{x} dx = \ln x + C$$

and so

$$y = \frac{\ln x + C}{x}.$$

Since  $y(1) = 2$  we have

$$2 = \frac{\ln 1 + C}{1} = C$$

Therefore the solution to the initial-value problem is

$$y = \frac{\ln x + 2}{x}.$$

## 2.5 Bernoulli Equations

### Definition 2.6

The differential equation

$$y' + a(x)y = b(x)y^n, \quad n \neq 0, \quad n \neq 1 \dots \dots \dots ( 5 )$$

where  $a$  and  $b$  are continuous functions on some interval  $I$ , is called a Bernoulli equation. To solve ( 5 ), multiply the equation by  $y^{-n}$  to obtain

$$y^{-n}y' + a(x)y^{1-n} = b(x)y^n.$$

The substitution  $z = y^{1-n}$ ,  $z' = (1 - n)y^{-n}y'$  transforms (5) into the following linear equation in  $z$  and  $x$ :

$$\frac{1}{1 - n} z' + a(x)z = b(x)$$

or

$$z' + (1 - n)a(x)z = (1 - n)b(x).$$

### Example 2.8

Solve the Bernoulli equation  $xy' + y = y^2 \ln x$ , For  $x > 0$ .

In this example  $a(x) = \frac{1}{x}$ ,  $b(x) = \frac{1}{x} \ln x$ , and  $n = 2$ . Therefore, we put  $z = y^{1-n} = \frac{1}{y}$ .

Then,  $z' = -\frac{1}{y^2}y'$ .

Inserting  $z = \frac{1}{y}$  and  $z' = -\frac{1}{y}y'$  into the differential equation, we get

$$z' - \frac{1}{x}z = -\frac{1}{x} \ln x$$

Thus, the resulting equation is a linear first order differential equation.

It can be solved using the integrating factor,

$$v(x) = e^{\int a(x)dx} = e^{-\int \frac{1}{x} dx} = \frac{1}{x}$$

Multiplying the differential equation by the integrating factor, we have

$$\frac{1}{x} z' - \frac{1}{x^2} z = -\frac{1}{x^2} \ln x$$

or

$$\left(\frac{z}{x}\right)' = -\frac{1}{x^2} \ln x$$

Integrating, we obtain

$$\begin{aligned}
\frac{z}{x} &= - \int \frac{1}{x^2} \ln x \, dx \quad (\text{By integrating by parts}) \\
&= \frac{1}{x} \ln x + \int \frac{dx}{x^2} \\
&= \frac{1}{x} \ln x + \int \frac{dx}{x^2} \\
&= \frac{1}{x} \ln x - \frac{1}{x} + C.
\end{aligned}$$

Multiplying by  $x$ , we have  $z = \ln x - 1 + Cx$ . Since  $z = \frac{1}{y}$ , the general solution to the problem is

$$y = \frac{1}{\ln x - 1 + Cx}.$$

### Example 2.9

Find every solution of the equation  $xy' = 3y + x^5y^{\frac{1}{3}}$ .

#### Solution

Rewrite the differential equation as

$$y' = \frac{3}{x}y + x^4y^{\frac{1}{3}}.$$

This is a Bernoulli equation for  $n = \frac{1}{3}$ . Divide the equation by  $y^{\frac{1}{3}}$

$$y^{-\frac{1}{3}}y' = \frac{3}{x}y^{\frac{2}{3}} + x^4.$$

Define the new unknown function  $z = y^{\frac{2}{3}}$ , calculate its derivative,  $z' = \frac{2}{3}y^{-\frac{1}{3}}y'$ , and introduce them in the differential equation,

$$\frac{3}{2}z' = \frac{3}{x}z + x^4 \quad \text{or} \quad z' - \frac{2}{x}z = \frac{2}{3}x^4$$

This is a linear equation for  $z$ . Integrate this equation using the integrating factor method. Then, the integrating factor is

$$v(x) = e^{\int a(x)dx} = e^{-\int \frac{2}{x}dx} = \frac{1}{x^2}$$

Therefore, the equation for  $z$  can be written as,

$$\frac{1}{x^2}z' - \frac{2}{x^3}z = \frac{2}{3}x^2$$

or

$$\frac{d}{dx} \left( \frac{1}{x^2}z \right) = \frac{2}{3}x^2.$$

Integrating, we obtain

$$\frac{1}{x^2}z = \frac{2}{3} \int x^2 \, dx = \frac{2}{9}x^3 + C$$

and the solution of the differential equation is

$$z = \frac{2}{9}x^5 + Cx^2$$

Once  $z$  is known we compute the original unknown  $y = \pm z^{\frac{3}{2}}$ , where the double sign is related to taking the square root. We finally obtain

$$y = \pm \left( \frac{2}{9}x^5 + Cx^2 \right)^{\frac{3}{2}}.$$